



Universität zu
Köln



Forschungszentrum Jülich



Tbilisi State
University

Measurement of Nucleon-Nucleon Elastic Scattering at Small Angles using the ANKE spectrometer

PhD Thesis by
Zara Bagdasarian

Supervisors: Dr. Dr. h.c. Hans Ströher (Universität zu Köln, FZJ)
Dr. Nodar Lomidze (Tbilisi State University, HEPI)

A thesis submitted in partial fulfillment for the
PhD Degree in Physics

under the joint supervision (Cotutelle) from University of Cologne
and Ivane Javakhishvili Tbilisi State University

Cologne/Tbilisi 2016

Measurement of Nucleon-Nucleon Elastic Scattering at Small Angles using the ANKE spectrometer

Inaugural-Dissertation

zur

Erlangung des Doktorgrades

der Mathematisch-Naturwissenschaftlichen Fakultät

der Universität zu Köln

(im gemeinsamen Promotionsverfahren (Cotutelle) mit der
Staatlichen Iwane-Dschawachischwili-Universität Tiflis)

vorgelegt von

Zara Bagdasarian

aus Tiflis

Köln/Tiflis 2016

Berichterstatter:

Prof. Dr. Dr. h.c. Hans Ströher,

Dr. Nodar Lomidze,

Prof. Dr. Merab Eliashvili

Tag der mündlichen Prüfung: 11.04.2016

ივანე ჯავახიშვილის სახელობის თბილისის სახელმწიფო უნივერსიტეტი
ზუსტ და საბუნებისმეტყველო მეცნიერებათა ფაკულტეტი
სადოქტორო პროგრამა: “ფიზიკა”

მცირე კუთხეებზე ნუკლონ-ნუკლონური დრეკადი გაბნევის შესწავლა ANKE სპექტრომეტრის გამოყენებით

ზარა ბაღდასარიანი

სადოქტორო დისერტაცია შესრულებულია

ფიზიკის დოქტორის
აკადემიური ხარისხის მოსაპოვებლად

თბილისის სახელმწიფო უნივერსიტეტს და კიოლნის უნივერსიტეტს (გერმანია)
შორის ერთობლივი ხელმძღვანელობის (Cotutelle) შესახებ ხელშეკრულების
ფარგლებში

დისერტაციის ხელმძღვანელები:

ჰანს შტროერი – კიოლნის უნივერსიტეტის პროფესორი,
იულიხის კვლევითი ცენტრის ბირთვული
ფიზიკის ინსტიტუტის დირექტორი

ნოდარ ლომიძე – ფიზიკა-მათემატიკის მეცნიერებათა დოქტორი,
თსუ მაღალი ენერგიების ფიზიკის ინსტიტუტის
უფროსი მეცნიერ თანამშრომელი

კიოლნი/თბილისი 2016

სადოქტრო პროგრამის ხელმძღვანელები:

მერაბ ელიაშვილი - თბილისის სახელმწიფო უნივერსიტეტის პროფესორი

მირიან ტაბიძე - ფიზიკა-მათემატიკურ მეცნიერებათა კანდიდატი, თსუ
მონაცემთა მოდელური ანალიზის ლაბორატორიის გამგე

ნორაირ აკოფოვი - ერევნის ფიზიკის ინსტიტუტის წამყვანი მეცნიერ-
თანამშრომელი

Abstract

A fundamental understanding of the nucleon-nucleon (NN) interaction is one of the ultimate goals of nuclear and hadron physics. Apart from their intrinsic importance for the study of nuclear forces, NN elastic scattering data are necessary, for example, in the modeling of meson production and other nuclear reactions at intermediate energies.

Quantum chromodynamics, the theory of strong interactions, is not able to determine the NN interaction from first principles because of its non-perturbative nature at intermediate energies (the coupling constants are too strong). Hence, phenomenological approaches are necessary to describe the NN interaction. Partial wave analyses (PWA), such as the ones regularly performed by the SAID (Scattering Analysis Interactive Dial-in) group, have proved to be truly invaluable tools over many years for researchers in this area. These analyses rely on the quantity and quality of the experimental measurements of various proton-proton (pp) and proton-neutron (pn) scattering observables at different energies over the full angular ranges.

The goal of many experiments conducted at Cooler Synchrotron (COSY) has been to provide PWA with the precision measurements of NN observables that are essential for these analyses. The experiments presented in this thesis have been carried out within the ANKE collaboration at COSY-Jülich, Germany. Data were obtained using polarised or unpolarised proton beams with kinetic energies from 0.8 GeV up to 2.8 GeV and unpolarised hydrogen or deuterium cluster-jet targets. The detection system of the ANKE spectrometer is the ideal set-up for carrying out refined measurements at the small scattering angles that had not previously been investigated.

The thesis comprises the measurements of the analysing power A_y and differential cross section $d\sigma/d\Omega$ in pp elastic scattering, and preliminary results for the analysing power A_y in the pn quasi-free elastic scattering. These new data close an important gap in the NN database at scattering angles up to 30° and energies up to 2.8 GeV.

The results obtained are compared to the predictions from the SAID PWA published in 2007 that were based on data from earlier experiments. The impact of the present results is demonstrated by the significant changes in the low partial waves of the updated SAID PWA, which includes the new ANKE measurements as well as the existing global data.

Zusammenfassung

Ein fundamentales Verständnis der Nukleon-Nukleon (NN) Wechselwirkung ist eines der wichtigsten Ziele der Kern- und Hadronenphysik. Neben ihrer zentralen Bedeutung für die Erforschung der Kernkräfte, sind Daten für elastische NN Streuung erforderlich, zum Beispiel bei der Modellierung von Mesonproduktion und anderen Kernreaktionen bei mittleren Energien.

Die Theorie der starken Wechselwirkung, Quantenchromodynamik, ist nicht in der Lage, die NN Wechselwirkung *ab initio* zu bestimmen, wegen der nichtstörungstheoretischen Natur bei mittleren Energien (die Kopplungskonstante ist zu stark). Daher sind phänomenologische Ansätze notwendig, die NN Wechselwirkung zu beschreiben. Partialwellenanalysen (PWA), z.B. jene, die regelmäßig von der SAID (Scattering Analysis Interactive Dial-in) Gruppe durchgeführt werden, haben sich als wirklich wertvolle Werkzeuge für Forscher in diesem Bereich erwiesen. Eine solche Analyse hängt stark ab von der Quantität und Qualität der experimentellen Messungen verschiedener Observablen von Proton-Proton (pp) und Proton-Neutron (pn) Streuung bei verschiedenen Energien und über den gesamten Winkelbereich.

Das Ziel vieler Experimente, die am Cooler Synchrotron (COSY) durchgeführt werden, ist, PWA mit den grundlegenden Präzisionsmessungen von NN Observablen zur Verfügung zu stellen. Die Experimente dieser Arbeit wurden an COSY-Jülich von der ANKE Kollaboration durchgeführt. Für die Messungen wurden polarisierte oder unpolarisierte Protonenstrahlen und unpolarisierter Wasserstoff oder Deuterium als Clusterjet-Target benutzt. Das Detektionssystem des ANKE-Spektrometer ist die ideale Einrichtung für präzise Messungen bei kleinen Streuwinkeln, die bisher nicht untersucht worden sind.

Die Dissertation beinhaltet die Messungen der Analysierstärke A_y und differentiellen Wirkungsquerschnitte $d\sigma/d\Omega$ in der pp elastischen Streuung und die vorläufigen Ergebnisse für die Analysierstärke A_y in pn quasi-freier elastischer Streuung. Diese neuen genauen Daten schließen eine wichtige Lücke in der NN Datenbank bei kleinen Streuwinkeln bis zu 30° und Energien bis zu 2.8 GeV.

Die erhaltenen Ergebnisse werden verglichen mit den SAID 2007 PWA, die auf Daten aus früheren Experimenten beruhen. Die Auswirkungen der vorliegenden Ergebnisse in Gestalt von signifikanten Veränderungen in den niedrigen Partialwellen der aktualisierten SAID PWA werden demonstriert.

ანოტაცია

ნუკლონ-ნუკლონური (NN) ურთიერთქმედების ფუნდამენტური შესწავლა არის ბირთვული და ადრონული ფიზიკის ერთერთი მთავარი ამოცანა. ბირთვული ძალების შესწავლის თავისთავადი მნიშვნელობის გარდა, NN დრეკადი გაზნევის მახასიათებლების ცოდნა აუცილებელია, მაგალითად, მეზონების დაბადების და სხვა ბირთვული რეაქციების მოდელირებისთვის შუალედურ ენერგიებზე.

ძლიერი ურთიერთქმედების თეორია - კვანტური ქრომოდინამიკა ძირითად პრინციპებზე დაყრდნობით ვერ აღწერს NN ურთიერთქმედებას შუალედურ ენერგიებზე, რადგან ბმის მუდმივის დიდი მნიშვნელობის გამო შეუძლებელია შემოთავაზების თეორიის გამოყენება. ამიტომ, NN ურთიერთქმედება ასეთ ენერგიებზე უნდა აღიწეროს ფენომენოლოგიური მიდგომით, კერძოდ, პროტონ-პროტონული (pp) და პროტონ-ნეიტრონული (pn) გაზნევის ამპლიტუდების პარციალურ ტალღებად გაშლის (ე.წ. ურთიერთქმედების ფაზური ანალიზის - PWA) საშუალებით. SAID (Scattering Analysis Interactive Dial-in) მონაცემთა ბაზა და ანალიზი წარმოადგენს ენერგიის ამ არეში ყველაზე სანდო ინფორმაციულ წყაროს, რომელიც იქმნებოდა ბოლო რამდენიმე ათეული წლის განმავლობაში. იგი დაფუძნებულია დღემდე არსებული pp და pn გაზნევის სხვადასხვა დამზერადი სიდიდეების ექსპერიმენტულ მონაცემებზე, სრულ კუთხურ ინტერვალში, სხვადასხვა შუალედურ ენერგიებზე.

COSY ამაჩქარებელზე ჩატარებული მრავალი ექსპერიმენტის მიზანს წარმოადგენდა მაღალი სიზუსტის ექსპერიმენტული მონაცემების შეგროვება PWA ანალიზისათვის. დისერტაციაში წარმოდგენილი ექსპერიმენტები ჩატარდა ANKE კოლაბორაციის მიერ არაპოლარიზებული ან პოლარიზებული პროტონული ნაკადისა და არაპოლარიზებული წყალბადის ან დეიტერიუმის ჭავლური სამიზნის გამოყენებით. ANKE სპექტრომეტრის დეტექტორების სისტემა არის იდეალური დანადგარი გაზნევის მცირე კუთხეების არეში დამზერადი სიდიდეების გაზომვებისთვის, სადაც დღემდე არ არსებობდა ექსპერიმენტული მონაცემები.

დისერტაციაში მოყვანილია pp დრეკადი გაზნევის რეაქციის ანალიზური უნარისა და დიფერენციალური კვეთის, pn კვაზიდრეკადი გაზნევის რეაქციის ანალიზური უნარის ექსპერიმენტული გაზომვის შედეგები. მიღებული შედეგები მნიშვნელოვნად ავსებენ NN მონაცემთა ბაზას გაზნევის მცირე (30° -მდე) კუთხეებსა და 2.8 GeV-მდე ენერგიის ინტერვალში.

დისერტაციაში მიღებული შედეგები შედარებულია SAID ანალიზის შედეგებთან, რომლებიც მიღებულ იქნენ 2007 წლამდე არსებულ ექსპერიმენტულ მონაცემებზე დაყრდნობით. ნაჩვენებია, აგრეთვე, ის ცვლილებები, რომლებიც SAID ანალიზზე მოახდინეს ANKE გაზომვების შედეგებმა დაბალი პარციალური ტალღების არეში.

Acknowledgements

The work presented in this thesis has been supported by the Forschungszentrum Jülich and Shota Rustaveli National Science Foundation (SFNSF) Fellowship and SRNSF Grant #DO/30/6-200/14.

First and foremost, I would like to thank my supervising Professor Dr. Dr. h.c. Hans Ströher, who manages to be a very humble and empathetic person regardless of the titles he has achieved. Thank you for always supporting me by giving me very rich advice that helped me grow during my PhD studies. Also, thanks for being so inspirational to me in the area of organization, whether for work, or purchasing tickets for my wedding, you're the most organized person I've ever met. And a very special thanks in regards to the Cotutelle agreement between the two universities, without your initiative, it would not have been possible.

I'm also sincerely grateful to my supervisor Dr. Nodar Lomidze for all of his help during these years of study. Thanks for teaching me how to question acquired knowledge and results in order to arrive at what is thought to be a consensus, and then to rethink the problem again. You have always given me exemplary guidance, which has in turn helped cultivate the necessary skills and ability to think independently as a graduate student.

I would like to gratefully acknowledge Dr. Andro Kacharava for his excellent guidance and cultivation of teamwork within the ANKE collaboration. Your substantial investment of time and energy that went into the building of the Georgian-German scientific bridge has created so many invaluable opportunities for students. One could write a separate thesis about everything that was achieved thanks to your exceptional enthusiasm and organizational skills.

A warm thanks to Professor Dr. Colin Wilkin, who not only possesses invaluable knowledge and insight into the field of physics, but also has a very charming personality. Your sage comments have always helped turn discussions and research in the right direction.

I would like to thank Dr. Ron Workman and Dr. Igor Strakovsky for showing such an interest in our results, and for collaborating with such efficiency. I sincerely hope to get the chance to thank both of you in person someday.

I want to thank Dr. David Mchedlishvili who always provides smart advice in incredible detail and depth. I cannot forget Dr. David Chiladze who was very helpful at the beginning of my PhD studies. I would also like to thank Dr. Mirian Tabidze for all the help in the university and working with students.

A great debt of gratitude is also owed to Dr. Giorgi Macharshvili, Dr. Sergey Dymov and Dr. Sergey Barsov. Thank you all for the close collaboration and the many fruitful discussions and explanations.

My heart is full of the warmest memories, which I have shared with so many amazing and kind people during last three years. I would like to mention Gisela

and Heinz-Günter Haubold who not only helped to tremendously improve my German skills, but can crash anybody's stereotypes about "cold Germans". I want to stay young like you when I grow up.

I wish to thank my friends who made the time in Jülich so memorable and fun. Thank you for all the interesting, thought-provoking conversations during lunch breaks and for the many laughs in the evenings. I would especially like to thank la Zhanna who has stayed my closest friend even after moving to the sunny south of France, as well as the best girls' lunch crew: Maria, Aude, Ellen and Oli.

I think one of my greatest fortunes is being blessed with a family that has supported me in all of my endeavors even when we live thousands of kilometers apart. I would like to thank my parents, who taught me that knowledge is the best treasure, you can gather during your whole life, and infected me with their love for math and curiosity for the world around us. You showed me how to do everything with great love and passion and how to create and keep strong family and friend bonds. I'm infinitely grateful to you for everything I have and everything I am today. I see traits of both of you in myself and I follow your guidance every day, even when time difference and busy schedule prevent me from calling on the daily basis. But you know, "vse xorosho".

A special thanks to the most awesome sister, who is also the kindest person on the planet. Thank you for all the support from you, Daniel and that bundle of awesomeness you two created, Kristen. I love you guys very very much, and I hope we get to spend more time together in the future #MoreVacation4Americans. I am also grateful to my new Italian family, who welcomed me so warmly, that my heart acquired another place to call home.

Finally, I would like to acknowledge Jülich for becoming the place where I met my wonderful husband Andrea. Thank you, Andrea, for all of the care and support you have given me, even at such a long distance. Thank you for being such a fun and comfortable person to be with. Here is to many years to come for the family of two PhDs exploring the world.

*To my wonderful parents,
Luiza and Amik,*

*and amazing hubby,
Andrea*

Contents

Abstract	vii
Zusammenfassung	ix
Abstract in Georgian	xi
Acknowledgements	xiii
Contents	xvii
Abbreviations	xxi
1 Introduction	1
1.1 Structure of matter	1
1.1.1 Structure of the nucleon: historical overview	3
1.2 Nucleon-nucleon scattering	5
1.2.1 EDDA's legacy	5
1.3 Synopsis of the thesis work	6
2 Formalism in polarised experiments	9
2.1 Polarisation formalism	9
2.1.1 Spin structure $1/2 + 1/2 \rightarrow 1/2 + 1/2$	10
2.1.2 Coordinate system	11
2.2 Spin observables	12
2.3 Cross-ratio method	13
3 Experimental setup	17
3.1 COSY facility	17
3.1.1 Polarised ion source at COSY	19
3.1.2 Depolarising resonances at COSY	20
3.2 EDDA polarimeter	22
3.3 ANKE spectrometer	24
3.3.1 Unpolarised cluster-jet target	24
3.3.2 Forward Detector (FD)	26

3.3.3	Silicon Tracking Telescopes (STT)	27
3.4	Experimental conditions	28
4	Analysing power in proton-proton elastic scattering	31
4.1	Beam polarisation measurement using EDDA	32
4.1.1	Beam polarisation uncertainty	34
4.2	Asymmetry measurement using ANKE	34
4.2.1	Event selection by STT	34
4.2.2	Asymmetry measurement using STT	36
4.2.3	Event selection by FD	37
4.2.4	Asymmetry measurement using FD	38
4.3	Sources of the systematic uncertainties	40
4.3.1	Difference in polarisation values for two polarisation modes	40
4.3.2	Polar angle accuracy	41
4.3.3	Detector efficiencies stability	42
4.3.3.1	Summary on systematic uncertainties	44
4.4	Results and discussion	44
4.5	Conclusion	47
5	Cross section in proton-proton elastic scattering	49
5.1	Luminosity determination	50
5.1.1	Beam intensity measurement	50
5.1.2	Target density determination	52
5.1.2.1	Schottky noise	53
5.1.2.2	Frequency shift	54
5.1.2.3	COSY rest gas	56
5.1.2.4	Momentum compaction factor	57
5.1.2.5	Luminosity summary	58
5.2	Event selection by FD	59
5.3	Systematic uncertainties	60
5.4	Results and discussion	60
5.4.1	Comparison with LNPI results at 1 GeV	62
5.4.2	Comparison with ANL results at 2.0 and 2.8 GeV	62
5.4.3	Impact on the SAID solution	63
5.5	Conclusion	67
6	Analysing power in proton-neutron quasi-free elastic scattering	69
6.1	Beam polarisation	70
6.2	Event selection	70
6.3	Asymmetry determination	72
6.3.1	Normalisation via proton-deuteron elastic scattering	73
6.4	Background correction	75
6.5	Validity of spectator model	76
6.5.1	Quasi-free elastic proton-proton scattering	77
6.6	Results and discussion	78

6.7 Conclusion	81
7 Summary and outlook	83
 Appendices	 87
A Angular ranges of EDDA semi-rings	89
B Numerical values of A_y in $\vec{p}p$ elastic scattering	91
C Numerical values of $d\sigma/d\Omega$ in pp elastic scattering	99
D Numerical values of A_y in $\vec{p}d$ elastic scattering	109
E Numerical values of A_y in $\vec{p}n$ quasi-free elastic scattering	113
 List of Figures	 115
List of Tables	121
Bibliography	125
 Erklärung	 137
Publikationen	138
Lebenslauf	141

Abbreviations

SM	S tandard M odel
PSA	P hase S hift A nalysis
PWA	P artial W ave A nalysis
QCD	Q uantum C hromo D ynamics
SAID	S cattering A nalysis I nteractive D ial-in
COSY	C Ooler S Ynchrotron
ANKE	A pparatus for S tudies of N ucleon and K aon E jectiles
EDDA	E xcitation Function D ata A cquisition Designed for A nalysis of Phase Shifts
STT	S ilicon T racking T elescope
FD	F orward D etector
MWPC	M ulti W ire P roportional C hamber
MWDC	M ulti W ire D rift C hamber
LEP	L ow E nergy P olarimeter
BCT	B eam C urrent T ransformer
RF	R adio F requency
PM	P hoto M ultiplier
ANL	A rgonne N ational L aboratory
LNPI	L eningrad N uclear P hysics I nstitute

Chapter 1

Introduction

What are the fundamental blocks of matter? How did the universe evolve just after the Big Bang? What are the mechanisms behind the various interactions? There is a wealth of fundamental questions with profound significance for our understanding of Nature and the structure of matter of which we and our universe are composed. Answering many of these questions lies at the heart of nuclear science.

One of the key moments for nuclear physics was the Geiger–Marsden experiment (also known as the Rutherford gold foil experiment) that showed how profoundly our understanding of the matter can be changed via a simple scattering experiment. Since then many increasingly sophisticated scattering experiments have been performed and we have learned a lot about the fundamental particles, but there are still many questions that we need to seek answer to.

The first two sections of this chapter will provide a short overview of the structure of matter, as we understand it today, and a short historical overview of our insight into the structure of the nucleon. The following sections describe the importance of the nucleon-nucleon scattering in general and the motivation of the experimental investigations, described in this dissertation.

1.1 Structure of matter

Our perception of the structure of matter has changed many times during the history and most rapidly since the end of the 19th century, when atoms were still thought to be the most basic, indivisible building blocks of matter, to the latter

half of the 20th century, when all known subatomic particles were organised within the Standard Model (SM).




FIGURE 1.1: The Standard Model of elementary particles (schematic depiction), with the three generations of matter in the first three columns, gauge bosons in the fourth column, and the Higgs boson in the fifth [1].

The Standard Model (SM) is the quantum field theory that describes all different kinds of interactions (except gravity, for which a quantum field theoretical description has not been achieved yet) and classifies the elementary particles. The elementary particles constituting the ordinary matter are the fundamental fermions (half-integer spin particles), namely quarks and leptons; the gauge bosons (integer-spin particles) mediate forces, while Higgs boson is responsible for the intrinsic mass of particles. The schematic depiction of SM elementary particles, with the three generations of matter, the Higgs and gauge bosons, is given in Figure 1.1.

The four fundamental forces are: the gravitational, electromagnetic, weak and strong forces. Even though weak and strong interactions are short ranged and hence were left unnoticed for a long time, nowadays we know their important role in the existence of matter. The weak force is involved in radioactivity, causing unstable atomic nuclei to decay, and plays important role in powering the sun's thermonuclear process. The strong interaction is responsible for holding quarks together, and consequently binding protons and neutrons into stable nuclei.

1.1.1 Structure of the nucleon: historical overview

Nucleons (protons and neutrons) are the lowest-energy bound states of quarks and gluons. They represent the simplest form of observable matter and comprise more than 99% of the mass of the visible universe. Most of these nucleons are in the core of atoms, the same atoms that everything we see on the daily basis is made from, including ourselves. So it comes as no surprise that nucleons have been studied with such a scrutiny during the last century.

Rutherford has started to use the word “proton” for the hydrogen nuclei after the first reported nuclear reaction $^{14}\text{N} + \alpha \rightarrow ^{17}\text{O} + p$ was observed and protons were identified as part of all nuclei. However, it was only after the discovery of neutron by James Chadwick in 1932 that the basic structure of nuclei and nuclear isotopes could be understood. Nevertheless, the nucleons were thought to be elementary particles not for long: already in 1933 first glimpse of an internal structure of the nucleon was observed, when magnetic moments of protons and neutrons were measured. In 1964 Gell-Mann [2] and, independently, Zweig [3] proposed a theory that nucleons are composed of point like particles called quarks. These quarks were postulated to have spin $-1/2$, a fractional electric charge, and came in different types called flavours. Soon after the electron scattering experiment that showed that nucleons are indeed composed of quarks took place at the Stanford Linear Accelerator Centre (SLAC) [4, 5]. Combinations of different flavours of quarks comprise baryons (built up from three quarks) and mesons (a quark and an anti-quark). These two groups of particles are categorized as hadrons.

Quantum Chromodynamics (QCD), the gauge field theory of the strong interaction, describes the interaction of quarks through the exchange of massless vector gauge bosons, the gluons. QCD follows the formalism of Quantum Electrodynamics (QED), which has a coupling constant a that describes the strength of the electromagnetic interaction. The underlying $\text{SU}(3)$ gauge structure, rather than the simple $\text{U}(1)$ of QED, implies many analogies, but also basic new features. The carriers of the strong force are 8 massless gluons in analogy to the photon for the electromagnetic force. An important new aspect is that the gluons, carrying a new quantum number called colour, can interact with each other.

As a consequence of the gluon self-coupling, QCD implies that the coupling strength α_s , the analogue to the fine structure constant α in QED, becomes large at large distances, or equivalently, at low momentum transfers. Therefore, QCD provides a qualitative reason for the observation that quarks do not appear as free particles

but only exist as bound states of quarks, forming hadrons. Quarks carry one out of three different colour charges: red, blue and green, while hadrons are colourless bound states. The gluons also carry the colour charge and therefore can interact with each other.

QCD does not predict the actual value of α_s . For large momentum transfers Q , however, it determines the functional form of the energy dependence of α_s , illustrated in Figure 1.2. While an increasingly large coupling at small energy scales leads to the *confinement* of quarks and gluons inside hadrons, the coupling becomes small at high-energy or short-distance reactions; quarks and gluons are said to be *asymptotically free* [6].




FIGURE 1.2: The value of the "running" coupling constant α_s , as a function of the energy scale E . The curve that slopes downwards is a prediction of the asymptotic freedom in QCD, while the empty circles show the measurements that have been made [7].

At high energies a perturbative approach can be taken in the mathematical description of the interactions, with α_s as the expansion parameter. This kinematic regime is called perturbative QCD or just pQCD, and theoretical predictions can be well tested by experiments. However at lower energies the coupling constant becomes larger, and perturbative calculations no longer work. Therefore the force description in this energy range relies on the phenomenological approach. The experimental data of high quality and precision at low and intermediate energy are necessary for the full understanding of the strong interaction.

1.2 Nucleon-nucleon scattering

The nucleon-nucleon (NN) interaction is the prototype for the action of the nuclear forces. Data on NN scattering are necessary ingredients, not only for the understanding of nuclear forces, but also for the description of meson production and other nuclear reactions at intermediate energies.

The scattering amplitudes for the complete description of the NN interaction can be reconstructed from the phase shift analysis (PSA). It has proved to be a truly invaluable tool over many years for researchers working in this area. For an interpretation of the results obtained in this work, they will be compared to the most recently published [8] and modified calculations from the SAID (Scattering Analysis Interactive Dial-in) partial wave analysis (PWA) [9, 10]. The SAID facility is based at George Washington University, Washington DC, USA. It maintains a database which contains the world data on NN scattering among other reactions, and provides predictions from PWA of the data. Such an analysis is based on the measurement of various NN scattering observables at different energies over the full angular range.

Unpolarised experiments provide information only about the averaged spin effects. This means that a certain amount of information is being lost. Since strong interaction is spin-dependent, it is crucial to conduct polarised experiments to separate the spin-specific parts of interaction. The polarised experiments provide additional information on reaction mechanisms, indispensable for the partial wave analysis.

1.2.1 EDDA's legacy

It should be pointed out that the COSY-EDDA collaboration (Excitation function Data acquisition Designed for Analysis of phase shifts) have provided a perfect example of how polarised data have completely revolutionised partial wave analysis.

The data on the differential cross section in pp elastic scattering [11] were taken in a continuous ramp of the proton beam energy from 0.24 to 2.58 GeV using the CH_2 fibre target. Prior to the EDDA measurements, SAID solution was only valid up to 1.6 GeV. With more than 2000 points EDDA data completely dominated the SAID database above about 1 GeV.

EDDA collaboration has also made contributions in spin-dependent measurements: the data points were produced for the proton analysing power using the unpolarised

beam between 0.44 and 2.49 GeV incident on a polarised hydrogen target [12, 13]. In addition, pp spin correlations were studied in the same energy range [14]. However, due to the design of the EDDA detector, the experiments only extended over the central region of centre-of-mass (c.m.) angles, $30^\circ \lesssim \theta_{cm} \lesssim 150^\circ$.

1.3 Synopsis of the thesis work

Many accelerators around the world included the NN study into their research program, however even after many years of studies, there are still many gaps in our knowledge. As one can see in the Figure 1.3, even in the data base of the most basic reaction of proton-proton elastic scattering, there has been a significant gap at the small angles ($\theta_{cm} < 30^\circ$) above 1 GeV. The situation is much worse for the isoscalar $I = 0$ case of proton-neutron scattering (Figure 1.4).




FIGURE 1.3: Abundance plots of c.m. scattering angle (θ_{cm}) versus beam kinetic energy (T_{lab}) for experiments on the analysing power A_y (left) and for cross-section $d\sigma/d\Omega$ (right) in proton-proton elastic scattering [15].

The precision data at small angles have a potential to significantly influence PSA. Adding to the NN scattering data base was one of the major priorities of the ANKE collaboration. This thesis comprises data gathered over three ANKE experiments, dedicated to the understanding of NN scattering:

- the proton-proton elastic scattering studies
 - analysing power A_y in proton-proton elastic scattering using unpolarised hydrogen cluster target and polarised proton beam at six energies between 0.796 and 2.4 GeV;




FIGURE 1.4: Abundance plot of c.m. scattering angle (θ_{cm}) versus beam kinetic energy (T_{lab}) for experiments on the analysing power A_y in proton-neutron scattering [15].

- unpolarised differential cross sections $d\sigma/d\Omega$ at eight beam kinetic energies between 1.0 and 2.8 GeV.
- The proton-neutron quasi-elastic scattering study
 - analysing power A_y using unpolarised deuterium cluster target and polarised proton beam at six energies between 0.796 and 2.4 GeV.

While high-quality pp data from ANKE close a very important gap at small angles, proton-neutron (pn) data are a crucial contribution to the very incomplete pn data base.

The outline of the remainder of this thesis is as follows: Chapter 2 describes the common formalism of the polarised experiments, on which the following chapters will be based. In Chapter 3 the experimental setup for the ANKE experiments is introduced. Chapter 4 gives a detailed description of the data analysis and obtained results for the analysing power in $\vec{p}p$ elastic scattering. In Chapter 5 the specifics of the cross section measurements in pp elastic scattering are described. Chapter 6 presents the data analysis performed on proton-neutron quasi elastic scattering and preliminary results. The concluding remarks are presented in Chapter 7.

Chapter 2

Formalism in polarised experiments

Polarisation physics represents the section of physics devoted to investigate the statistical and dynamical characteristics of processes associated with spin, which is one of the fundamental characteristics of elementary particles and nuclei. The spin is a tool to investigate and test fundamental questions. Performing the unpolarised cross section measurements for non-zero spin particles implies that in the description and experimentally there is summing over initial-channel and averaging over final-channel spin states. This means that a certain amount of information is being lost.

In this chapter a short overview of the formalism necessary to investigate the *spin-specific* parts of nucleon-nucleon (NN) interaction is given. Namely, important aspects of polarised experiments necessary to extract the spin observables, and a method to reduce the systematic errors. Within this thesis, only interactions of $1/2$ spin particles are studied, hence only this simple case of $1/2$ spin particle scattering is described in the following.

2.1 Polarisation formalism

Quantum mechanics deals with statistical statements about the result of measurements on an ensemble of states (particles, beams, targets). In other words: by giving an expectation value of operators it provides probability amplitudes for the result of a measurement on an ensemble.

There are two limiting cases. One is the pure state, that is our knowledge of the system is complete, e.g. when all members of an ensemble are in the same spin

state. A special case is the spin state of a single particle, which is always completely polarised. In general, our knowledge of a system is incomplete and can only be described by superposition of such pure states, weighted with the probability of their occurrence in this superposition. Such a state is called a mixed state. The appropriate and also practical description of such states is by using the density operator ρ :

$$\rho = \sum_i p_i |\psi^i\rangle \langle \psi^i|, \quad (2.1)$$

where p_i is the probability of finding the ensemble in a quantum mechanical state characterised by $|\psi^i\rangle$. A pure state is represented by the density matrix:

$$\rho = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (2.2)$$

A completely unpolarised beam with all spin substates equally occupied has the density matrix

$$\rho = 1/2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 1/2 \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right]. \quad (2.3)$$

This corresponds to superposition of pure states with equal weights of 1/2. A general beam can be interpreted as a superposition of the two pure states defined with respect to the quantization axis with the contributions N_+ and N_- :

$$\rho = N_+ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + N_- \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} N_+ & 0 \\ 0 & N_- \end{pmatrix}. \quad (2.4)$$

The vector polarisation of a spin 1/2 system has form:

$$P = \frac{N_+ - N_-}{N_+ + N_-}. \quad (2.5)$$

2.1.1 Spin structure $1/2 + 1/2 \rightarrow 1/2 + 1/2$

Formalism of elastic scattering of the systems with spin structure $1/2 + 1/2 \rightarrow 1/2 + 1/2$, including the NN scattering, is described in detail in References [16] and [17]. In principle, there are 255 possible polarisation observables for this spin system + unpolarised differential cross section. However, for elastic scattering, parity conservation and time-reversal invariance will reduce this number to 25 for identical particles, for example pp scattering, and to 36 linearly independent

experiments for non-identical particles, such as pn scattering. Nucleon-nucleon scattering matrix M is presented as

$$M(k_f, k_i) = \frac{1}{2} \{ (a + b) + (a - b)(\sigma_1, \mathbf{n})(\sigma_2, \mathbf{n}) + (c + d)(\sigma_1, \mathbf{m})(\sigma_2, \mathbf{m}) + (\mathbf{c} - \mathbf{d})(\sigma_1, \mathbf{l})(\sigma_2, \mathbf{l}) + \mathbf{e}(\sigma_1 + \sigma_2, \mathbf{n}) \}, \quad (2.6)$$

here the amplitudes a, b, c, d , and e are complex functions of two variables: energy in c.m. and the scattering angle. $\mathbf{l}, \mathbf{m}$, and $\mathbf{n}$ are the c.m. basis vectors:

$$\mathbf{l} = \frac{\mathbf{k}_f + \mathbf{k}_i}{|\mathbf{k}_f + \mathbf{k}_i|}, \mathbf{m} = \frac{\mathbf{k}_f - \mathbf{k}_i}{|\mathbf{k}_f - \mathbf{k}_i|}, \mathbf{n} = \frac{\mathbf{k}_f \times \mathbf{k}_i}{|\mathbf{k}_f \times \mathbf{k}_i|}. \quad (2.7)$$

whereas σ_1 and σ_2 are Pauli matrices. After considering parity conservation, time-reversal invariance and the Pauli principle, there are $N = 5$ invariant, independent complex amplitudes in pp scattering, and $N = 6$ in np scattering. Thus in a complete experiment $2N - 1$ real quantities have to be measured by at least as many independent experiments: 9 for pp and 11 for np [18].

2.1.2 Coordinate system

For polarised experiments the appropriate definition of coordinate system is very important. In unpolarised reactions incoming and outgoing particle momenta define the scattering plane. A polarisation vector adds another direction, introducing an azimuthal dependence of observables. Suitable coordinate systems have been agreed upon in two international conventions on polarisation phenomena in nuclear reactions, in Basel [19] and in Madison [20]. A cartesian coordinate system is formed with z along the incident beam momentum $\mathbf{k}_i$, y along $\mathbf{k}_i \times \mathbf{k}_f$ where $\mathbf{k}_f$ is the scattered particle momentum, and x such as to define a right-handed coordinate system. Let's define $\mathbf{i}, \mathbf{j}$ and $\mathbf{k}$ unit vectors pointing along the x, y and z coordinate axes respectively. So, $\mathbf{k} = \mathbf{k}_i/k_i$, $\mathbf{j} \parallel (\mathbf{k}_i \times \mathbf{k}_f)$, and $\mathbf{i} \parallel (\mathbf{k} \times \mathbf{j})$.

The unit vector pointing along the spin quantization axis is denoted by $\mathbf{S}$; its direction is defined in terms of β , the angle between $\mathbf{S}$ and beam direction, and ϕ , the angle between its projection on xy plane and the y axis. In this system, the scattering is always in the xz plane, and the momentum vector of the scattered particle lies in the xz half-plane with positive x . The so-called “up” direction is defined by the transverse component of the spin quantization axis: $\mathbf{S}_\perp = \mathbf{S} - (\mathbf{S} \cdot \mathbf{k})\mathbf{k}$. Hence, according to an observer who is looking along the beam direction and is aligned




FIGURE 2.1: Madison convention for the definition of the coordinate system in the polarised experiments. The z axis is along the incident beam momentum, while the scattering is in the xz plane. The spin quantization axis $\mathbf{S}$ is defined in terms of β , the angle between $\mathbf{S}$ and beam direction, z -axis, and ϕ , the angle between its projection on xy plane and the y axis.

with the “up” direction, the scattering (positive x half-plane of xz plane) is to the left, if the y axis is along $\mathbf{S}_\perp$ ($\phi = 0^\circ$). Correspondingly, if $\phi = 180^\circ$, $\phi = 270^\circ$ and $\phi = 90^\circ$, then the scattering is to the right, up and down, respectively.

2.2 Spin observables

Although the spin observables depend only on a polar angle, the cross section including these observables generally exhibit also a dependence on the azimuthal angle. This dependence enters via the need to introduce a coordinate system in which the detector positions, as well as the polarisation direction, have to be described. The cross section for a polarised beam of spin 1/2 particles is

$$\sigma(\theta, \phi) = \sigma_0(\theta)[1 + P_y A_y(\theta)], \quad (2.8)$$

where $\sigma_0(\theta)$ is the cross section for the scattering of an unpolarised beam at the scattering angle θ , $A_y(\theta)$ is the analysing power of the reaction at the same angle and P_y is the y component of the beam polarisation:

$$P_y = \mathbf{P} \cdot \mathbf{j} \equiv \mathbf{P} \sin \beta \cos \phi \equiv \mathbf{P}_\perp \cos \phi, \quad (2.9)$$

where $\mathbf{P}$ is the beam polarisation, $\mathbf{j}$ is unit vector along the y axis, and $\mathbf{P}_\perp$ is the component of beam polarisation perpendicular to its direction of motion. Since in

our experiment we have a transversely polarised beam, we assume $\beta \approx 90^\circ$. Hence the difference between $P_\perp$ and P is neglected, and we set $P_\perp \equiv P$.

The Madison convention implies that for spin 1/2 particles the polarisation should be counted positive in the direction $(\mathbf{k}_i \times \mathbf{k}_f)$. Assuming a positive analysing power, this positive polarisation yields a positive left-right (L-R) asymmetry.

2.3 Cross-ratio method

Many sources of systematic uncertainties in the determination of the analysing power A_y can be neglected in the first order in case of left-right symmetrical arrangement of the detection system [21]. Let us consider symmetric two detector system, depicted in Figure 2.2. The actual number of counts recorded in a detector is

$$N(\theta, \phi) = nN_t\Omega E\sigma(\theta, \phi), \quad (2.10)$$

where n is the number of particles incident on the target, N_t is the number of target nuclei per cubic centimetre, Ω is a geometrical factor, defined by the detector, i.e. the solid angle subtended by the detector, and E is the detector efficiency. It is allowed that the solid angle factor, as well as the efficiency of detector 1 are different from those of detector 2. Therefore, Equation 2.10 for each detector will look like

$$N_1(\theta, \phi) = nN_t\Omega_1 E_1 \sigma_0(\theta) [1 + P_y A_y(\theta) \cos\phi], \quad (2.11)$$

$$N_2(\theta, \phi) = nN_t\Omega_2 E_2 \sigma_0(\theta) [1 + P_y A_y(\theta) \cos\phi]. \quad (2.12)$$

In Figure 2.2 beam polarisation direction "up" is depicted coming out of paper in red. In this case, detector 1 detects particles, that are scattered to the "left" ($\phi = 0^\circ$) and detector 2 gets particles that are scattered to the "right" ($\phi = 180^\circ$). Therefore,

$$N_1(\theta, 0) \equiv N_1^\uparrow \equiv L_1 = nN_t\Omega_1 E_1 \sigma_0(\theta) [1 + P_y A_y(\theta)], \quad (2.13)$$

$$N_2(\theta, \pi) \equiv N_2^\uparrow \equiv R_2 = nN_t\Omega_2 E_2 \sigma_0(\theta) [1 - P_y A_y(\theta)]. \quad (2.14)$$

If we now "flip" the polarisation $\mathbf{P} \rightarrow -\mathbf{P}$, the spin direction will be going into the paper (depicted in green). This is referred to as beam polarisation "down", and in




FIGURE 2.2: The idealistic symmetric arrangement of the two-detector system.. The two cases of beam polarisation are shown: “up” in red and “down” in green. The corresponding counts of particles scattered to the detectors are described in text.

this case detector 1 will be on the right and detector 2 on the left side.

$$N_1(\theta, \pi) \equiv N_1^{\downarrow} \equiv R_1 = n' N_t' \Omega_1 E_1 \sigma_0(\theta) [1 - P_y A_y(\theta)], \quad (2.15)$$

$$N_2(\theta, 0) \equiv N_2^{\uparrow} \equiv L_2 = n' N_t' \Omega_1 E_2 \sigma_0(\theta) [1 + P_y A_y(\theta)]. \quad (2.16)$$

Primes are used to indicate that the integrated charge and the effective target thickness may be not the same for the two runs. We can form geometrical means of number of particles scattered to the left $L \equiv \sqrt{L_1 L_2}$ and particles scattered to the right $R \equiv \sqrt{R_1 R_2}$.

$$L = [nn' NN' \Omega_1 \Omega_2 E_1 E_2]^{\frac{1}{2}} \sigma_0 [1 + P A_y(\theta)], \quad (2.17)$$

$$R = [nn' NN' \Omega_1 \Omega_2 E_1 E_2]^{\frac{1}{2}} \sigma_0 [1 - P A_y(\theta)]. \quad (2.18)$$

We can solve for $P A_y(\theta)$ and get the left-right asymmetry ε

$$\varepsilon = \frac{L - R}{L + R} = \frac{2 P A_y [nn' NN' \Omega_1 \Omega_2 E_1 E_2]^{\frac{1}{2}} \sigma_0}{2 [nn' NN' \Omega_1 \Omega_2 E_1 E_2]^{\frac{1}{2}} \sigma_0} = P A_y(\theta), \quad (2.19)$$

which is independent of the relative detector efficiencies (E_1, E_2) , solid angles (Ω_1, Ω_2) , relative integrated charge (nn') and target thickness variations. (NN') . n and N , quantities common to the two channels, can be averaged over the data acquisition time (in one run), while E and Ω , quantities different in the two channels, must not vary with time [21]. We can define the geometric mean of the

number of particles detected by detector 1 in two runs as N_1 :

$$N_1 \equiv \sqrt{L_1 R_1} = \Omega_1 E_1 \sigma_0(\theta) N N' n n' [1 - (P A_y)^2]^{\frac{1}{2}}, \quad (2.20)$$

and for particles detected by detector 2 we have

$$N_2 \equiv \sqrt{L_2 R_2} = \Omega_2 E_2 \sigma_0(\theta) N N' n n' [1 - (P A_y)^2]^{\frac{1}{2}}. \quad (2.21)$$

Monitoring on the ratio

$$\frac{N_1}{N_2} = \frac{\Omega_1 E_1}{\Omega_2 E_2} \quad (2.22)$$

provides the check on the performance of the apparatus; this variable is required to be constant in time if the asymmetry determination is to be accurate. The statistical error associated with a measurement of the asymmetry ε is given by the means of geometrical means L and R :

$$\delta\varepsilon = \sqrt{\frac{1 - \varepsilon^2}{L + R}}. \quad (2.23)$$

Chapter 3

Experimental setup

All the results presented in this thesis, have been obtained using the data gathered at the ANKE spectrometer at COSY-Jülich. COSY facility with the polarised ion source are introduced in the Sections 3.1 and 3.1.1. Even though the COSY facility includes many experimental possibilities, this chapter includes the description of only those experimental equipments that have been actively used in the described experiments. Namely, the EDDA polarimeter and ANKE spectrometer are discussed in Sections 3.2 and 3.3 respectively. Finally, an overview of the different experimental conditions, under which the present data have been acquired, is given in Section 3.4.

3.1 COSY facility

The COSY accelerator and storage ring, shown schematically in Figure 3.1, serves the quests of the fundamental research in the Institute of Nuclear Physics of Jülich research centre (Forschungszentrum Jülich). The name COoler SYnchrotron (COSY) refers to the two phase space cooling mechanisms integrated in the ring. A beam with a momentum below 0.6 GeV/c can be cooled with the electron cooler [22, 23]. The stochastic cooler is used to manipulate the beam above a momentum of 1.5 GeV/c [24]. The acceleration process of the COSY beam consists of several stages. Negative ion sources can produce unpolarised and polarised hydrogen and deuterium ions, which are then accelerated by JULIC cyclotron up to 300 MeV/c for H^- and up to 600 MeV/c for D^- . These pre-accelerated ions are stripped off their electrons and the remaining protons or deuterons are injected in COSY ring with 183.4 m circumference, here particles can be accelerated and

stored at any momentum in the range from 300 MeV/ c to 3.65 GeV/ c [25]. At the injection beamline, the Low Energy Polarimeter (LEP) can provide a polarisation measurement of the beam generated by the polarised ion source.




FIGURE 3.1: The COSY accelerator facility at Forschungszentrum Jülich. The positions of the ANKE spectrometer and the EDDA polarimeter at the ring are shown. In the bottom of the figure the sources providing polarised and unpolarised protons (or deuterons), the cyclotron that accelerates the particles to the injection energy and the Low Energy (LE) Polarimeter can be seen.

Transversely polarised proton beams are available with intensities up to $1 \cdot 10^{10}$ particles with a polarisation up to 70%. For deuterons an intensity of $3 \cdot 10^{10}$ with vector and tensor polarisation of more than 70% and 50% were achieved respectively. The two 40 m long straight sections are designed to serve the internal experiments. Figure 3.1 demonstrates the COSY accelerator facility with the

positions of the ANKE spectrometer and the EDDA polarimeter, used during the experiments that are subject of this thesis.

3.1.1 Polarised ion source at COSY

The polarised ion source at COSY consists of three groups of components: the pulsed atomic beam source, the caesium beam source, and the charge-exchange and extraction region [26]. The schematic layout of the setup is shown in Figure 3.2. The use of atomic hydrogen allows one to take advantage of the large magnetic moment of the electron. The nuclear spin, in turn, is affected by its coupling to the electron [27].





FIGURE 3.2: Setup of the polarised ion source at COSY [28]. The negatively charged ions are produced in a charge exchange process between a neutral nuclear polarised hydrogen beam and a fast neutral cesium beam.

The neutral polarised hydrogen H^0 beam is produced in the atomic beam source, consisting of RF dissociator and a sextupole separation magnet. First, the gas molecules are dissociated in a RF discharge (300-400 W) and a high degree of dissociation is maintained by adding small amounts of nitrogen and oxygen that reduces surface and volume recombination. The atoms are cooled to about 30 K by passing through an aluminium nozzle of 20 mm length and 3 mm diameter. By slowing down the atoms, the acceptance of the hexapole system and dwell time in the charge-exchange are increased. The first sextupole magnet produces electron state polarisation by defocusing atoms with the electron spin state $m_j = -1/2$. Remaining beam of atoms with $m_j = +1/2$ is then focused by the second sextupole

magnet. The nuclear polarisation is provided by two RF transitions switching between the hyperfine substates of the hydrogen atoms [29].

Afterwards, the atomic $\vec{H}^0$ beam with now high nuclear polarisation collides with the fast neutral caesium (Cs) beam. Thanks to the significantly higher electronegativity, a hydrogen atom acquires an additional electron from a caesium atom, and becomes negatively charged, according to



The fast Cs^0 beam, needed for the abovementioned reaction, is produced in two steps. First, Cs vapor is thermally ionized on a hot (1200°) porous tungsten surface at an appropriate beam potential of about 40-60 kV, where the cross section for the charge-exchange reaction has its maximum. Second, the beam is focused by a quadrupole triplet to a neutraliser that consists of caesium oven, a cell filled with caesium vapour, and a magnetically driven flapper valve between the oven and the cell. The remaining Cs^+ beam is deflected in front of the solenoid into a Faraday cup, while the fast neutralised caesium atoms enter the charge-exchange region. A neutraliser efficiency is typically over 90%.

In the charge exchange region the nuclear polarisation is preserved by the longitudinal magnetic field. A small electrostatic gradient field guides the very slow H^- ions to the extraction orifice, where they are deflected by a 90°electrostatic toroidal deflector into the injection beamline of the cyclotron. In the final stage, a Wien filter separates the H^- ions from electrons and other background. The Wien filter is rotatable around the beam axis, providing any orientation of the polarisation vector. In order to avoid the polarisation loss during the acceleration, spin orientation parallel to that of cyclotron magnetic field is chosen[30].

3.1.2 Depolarising resonances at COSY

This section is dedicated to the description of the difficulties in the acceleration of the polarised proton beam. For an ideal planar closed-loop accelerator with a vertical guide field, the particle spin vector precesses around the vertical axis. In this way the vertical beam polarisation is preserved. The spin motion in an external electromagnetic field is governed by the Thomas–BMT equation, leading to a spin tune $\nu_{sp} = \gamma G$, which describes the number of spin precessions of the central beam per revolution in the ring. G is the anomalous magnetic moment of

the particle ($G = 1.7928$ for protons, -0.1423 for deuterons), and $\gamma = E/m$ is the Lorentz factor. During the acceleration of a vertically polarised beam, depolarising resonances are crossed if the precession frequency of the spin γG is equal to the frequency of the encountered spin-perturbing magnetic fields. In a strong-focusing synchrotron like COSY, two different types of strong depolarising resonances are excited, namely imperfection resonances caused by magnetic field errors and misalignments of the magnets, and intrinsic resonances excited by horizontal fields due to the vertical focusing [31].

In the momentum range of COSY, five imperfection resonances have to be crossed for protons. The existing correction dipoles of COSY are utilised to overcome all imperfection resonances by exciting adiabatic spin flips without polarisation losses. The number of intrinsic resonances depends on the superperiodicity of the lattice. The magnetic structure of COSY allows one to choose a superperiodicity of $P = 2$ or 6 . A tune-jump system consisting of two fast quadrupoles has been developed especially to handle intrinsic resonances at COSY [32].

The imperfection resonances for protons in the momentum range of COSY are listed in Table 3.1. They are crossed during acceleration, if the number of spin precessions per revolution of the particles in the ring is an integer ($\gamma G = k, k$ is integer). The resonance strength depends on the vertical closed orbit deviation.

γG	T_p GeV	p GeV/ c	y_{co}^{rms} mm	ϵ_r 10^{-3}	P_f/P_i
2	0.1084	0.4638	2.3	0.95	-1.00
3	0.6318	1.2587	1.8	0.61	-0.88
4	1.1551	1.8712	1.6	0.96	-1.00
5	1.6785	2.4426	1.6	0.90	-1.00
6	2.2018	2.9964	1.4	0.46	-0.58

TABLE 3.1: Resonance strength ϵ_r and the ratio of preserved polarisation P_f/P_i at imperfection resonances for a typical vertical orbit deviation y_{co}^{rms} , without considering synchrotron oscillation.

A spin flip occurs at all resonances if synchrotron oscillations are not considered. However, the influence of synchrotron oscillation during resonance crossing cannot be neglected. After the first imperfection resonance, the calculated polarisation with a momentum spread of $\Delta p/p = 1 \times 10^{-3}$ and a synchrotron frequency of $f_{syn} = 450$ Hz is about $P_f/P_i \approx -0.85$. The resonance strength of the first imperfection resonance has to be enhanced to $\epsilon_r = 1.6 \times 10^{-3}$ to excite spin flips with polarisation losses of less than 1%. At the other imperfection resonances the effect

of synchrotron oscillation is smaller, due to the lower momentum spread at higher energies. Vertical correction dipoles or a partial Siberian snake could be used to preserve polarisation at imperfection resonances by exciting adiabatic spin flips. Simulations indicate that an excitation of the vertical orbit with existing correction dipoles by 1 mrad is sufficient to adiabatically flip the spin at all imperfection resonances. In addition, the solenoids of the electron-cooler system inside COSY are available for use as a partial snake. They are able to rotate the spin around the longitudinal axis by about 8° at the maximum momentum of COSY. A rotation angle of less than 1° of the spin around the longitudinal axis already leads to a spin flip without polarisation losses at all five imperfection resonances [33]. The number of intrinsic resonances depends on the superperiodicity P of the lattice, which is given by the number of identical periods in the accelerator. The COSY ring consists of two 180° arc sections connected by straight sections. The straight sections can be tuned as telescopes with 1:1 imaging, giving a 2π betatron phase advance. In this case the straight sections are optically transparent and the arcs contribute to the strength of intrinsic resonances. One then obtains for the resonance condition $\gamma G = k \times P \pm (Q_y - 2)$, where k is an integer and Q_y is the vertical betatron tune. The magnetic structure in the arcs allows adjustment of the superperiodicity to $P = 2$ or 6 . The corresponding intrinsic resonances in the momentum range of COSY are listed in Table 3.2.

P	γG	T_p MeV	p MeV/ c	ϵ_r 10^{-3}
2	$6 - Q_y$	312.4	826.9	0.26
2	$0 + Q_y$	950.7	1639.3	0.21
2,6	$8 - Q_y$	1358.8	2096.5	1.57
2	$2 + Q_y$	1997.1	2781.2	0.53
2	$10 - Q_y$	2405.2	3208.9	0.25

TABLE 3.2: Resonance strength ϵ_r of intrinsic resonances for a normalized emittance of 1π mm mrad and vertical betatron tune of $Q_y = 3.61$ for different superperiodicities P .

3.2 EDDA polarimeter

Even though the polarisation of the beam is measured at the injection via Low Energy Polarimeter (LEP), it is important to control the polarisation of the beam after it goes through all the depolarisation resonances and reaches the flattop (constant kinetic energy). This was achieved via EDDA measurements during the last 20 seconds at the end of the every cycle.

The EDDA experiment was initially conceived to provide high precision pp elastic scattering data in the COSY energy range (0.5 - 2.5 GeV), but later has been modified to be used as the internal polarimeter. The EDDA-polarimeter is comprised of the $7\ \mu\text{m}$ diameter carbon fibre target and 2×29 semi-ring scintillators that intercept protons within the polar angle range from 11.1° to 42.7° in the laboratory coordinate system (Appendix A) [12].




FIGURE 3.3: The schematic layout of the full EDDA detector setup (left) and photo (right) of the EDDA polarimeter.

The outside layer consists of a series of rings that wrap around the bars and each intercepts a narrow range of scattering angles from the target. The stripped-down version of the EDDA detector used as a polarimeter at COSY does not include the bars. It was calibrated during the EDDA data-taking periods against the full detector setup. As demonstrated in Figure 3.3, the rings (R) are split into left and right halves. Each half is connected to a light guide and single photomultiplier tube. The triggers are generated for each of the solid semiring-shaped scintillators, signal from which were counted in the so-called scalars. The time-marking system uses a precise clock to provide time for each event trigger. These times are stored and passed to the event processing software. The scalar rates are then read out separately for each ring and the two spin directions, and making use of the cross-ratio method, and known ring effective analysing powers [34], the beam polarisation is calculated. See the details in Section 4.1.

3.3 ANKE spectrometer

The Apparatus for Studies of Nucleon and Kaon Ejectiles (ANKE) is an internal experiment in one of the straight section of COSY. It consists of the magnetic system (three dipole magnets), unpolarised hydrogen or deuteron cluster-jet target, and different detection systems, from which we have mostly used for this experiment only Forward Detector (FD) and Silicon Tracking Telescopes (STTs). In Figure 3.4 only those parts of the spectrometer are shown that are relevant for this experiment.




FIGURE 3.4: The ANKE spectrometer setup (top view), showing the positions of the hydrogen cluster-jet target, the silicon tracking telescopes (STT), and the forward detector (FD).

The main purpose of the ANKE magnetic system is to separate the ejectiles from the circulating COSY beam in order to identify them and analyse their momentum. The first dipole magnet D1 deflects the circulating beam by an angle α off its straight path onto the target; the spectrometer dipole magnet D2 (beam deflection -2α) separates the produced particles from the beam for momentum analysis; finally D3, identical to D1, leads the unscattered beam particles back onto the normal orbit [35]. The deflection angle of the beam can be adjusted to optimise the magnetic field up to 1.56 T independent of the COSY beam momentum.

3.3.1 Unpolarised cluster-jet target

For the ANKE experiments target with thickness of 10^{13} to 10^{15} *atoms/cm²* is typically used. For experiments that require unpolarised targets, it is provided by

cluster-jet target device, shown in Figure 3.5. It consists of three main parts: the cluster-jet source, the scattering chamber, and the cluster-jet beam dump [36].




FIGURE 3.5: The cluster-jet target installed at ANKE.

Figure 3.6 illustrates the schematic overview of the cluster production process. The hydrogen or deuterium gas at pressures of 18 bar is cooled down to temperatures of 20-30 K and pressed through a Laval nozzle. Adiabatic expansion of the cooled beam further cools down the beam and the oversaturated gas spontaneously condensates to clusters with typical size of $10^3 - 10^4$ atoms. Even though the chosen temperature, pressure and nozzle geometry have already been optimized, only a small part of used gas forms clusters. Hence the skimmer (an opening of $700 \mu\text{m}$) is used to separate the cluster jet from the surrounding gas. The final shape of the cluster jet is defined by a second opening, collimator. Finally, the cluster beam is separated from the residual gas by a skimmer.

The scattering (analysing) chamber is equipped with a scanning rod with a thickness of 1.0 mm which is controlled by a stepper-motor and can be positioned in




FIGURE 3.6: Cluster production process in a Laval-nozzle. The cluster-jet beam is extracted and shaped using the skimmer. Note the exaggerated scale, the opening of the skimmer is only $700 \mu\text{m}$.

units of $1/24 \text{ mm}$. When the rod is placed inside the cluster beam, a part of the beam is stopped and converted into a gas load which can be recorded by an ionization vacuum meter. In that way information on the cluster beam size and position can be obtained. Furthermore, if the rod is placed at a fixed position inside the cluster beam, this system allows the density of the beam to be monitored.

The part of the cluster-jet beam, which does not interact with the COSY beam, is collected in the beam dump. It consists of three cryopumps and one turbo molecular pump mounted at the end.

3.3.2 Forward Detector (FD)

Forward Detector (FD) is located in the gap of 1.6 m between the D2 and D3 dipole magnets. The closeness of the FD part to the COSY beam pipe introduces the requirement for the system to operate at rather high counting rates of 10^5 s^{-1} and more.

The FD comprises two multiwire proportional (MWPC) and one drift chamber (MWDC) as well as a two-plane scintillation hodoscope. The information from MWPC and MWDC chambers are used for track reconstruction at ANKE. The existence of the strong magnetic field of the D2 ensures a good spatial separation of tracks with different mass-to-charge (m/q) ratio. Using the hit information from different layers of the MWPC and MWDC and the geometrical position of the target, tracks are found from the overall fit procedure. Details on the track-finding algorithm and the track-reconstruction software can be found in [37] [38].

The MWPCs have four wire planes each, with two horizontally (X) and two vertically (Y) aligned wires as well as two-strip planes, which are inclined by $\pm 30^\circ$ with respect to the wires. High spatial resolution of less than 1 mm is required from the Multiwire Proportional Chambers (MWPCs), in order to achieve momentum resolution of about 1%, which is essential for distinguishing proton-proton pairs with low excitation energy.

The forward hodoscope is composed of two planes of vertically aligned scintillators from polystyrene. The first and second planes contain 8 and 9 scintillators, respectively. In each plane, counters that are placed close to the COSY beam pipe, have smaller thickness (15 mm) and width (varying between 40 and 60 mm), compared to those responsible for lower momentum region (20 mm thick, 80 mm wide). The height of all scintillators is 360 mm. Each of the scintillators is read out by two photomultiplier tubes placed on both ends. They provide timing as well as the amplitude signal. The timing signal can be used to form a trigger and also to measure the differences of the arrival times of particle pairs. A typical time resolution for events with two registered particles is around 0.5 ns. The amplitude signal from photomultipliers provides information about the energy loss in the scintillator, which can be measured with 10% accuracy [39].

3.3.3 Silicon Tracking Telescopes (STT)

Two Silicon Tracking Telescopes are placed inside the vacuum chamber in a ϕ -symmetric (left-right) arrangement close to the beam-target overlap (schematically shown in Figure 3.4). Each STT consists of three individual double-sided silicon strip detectors of different thickness. The basic configuration has a 65 μm first layer, a 300 μm second layer and to ensure stopping of protons with kinetic energy up to 40 MeV, a third layer of 5100 μm thickness. The first layer is placed 28 mm from the centre of the beam pipe. The distance between the two first layers is set to 20 mm. The photograph of the STT detectors, along with the cooling system and front-end read-out electronics, is given in Figure 3.7.

Measuring the energy losses in the individual layers of the telescope allows the identification of stopped particle by the $\Delta E/E$ method. From the Bethe-Bloch formula, it can be seen that the energy loss in matter depends on the charge ze and the velocity of the incident particle $v = \beta c$.

$$-\frac{dE}{dx} = \frac{4\pi}{m_e c^2} \frac{nz^2}{\beta^2} \left(\frac{e^2}{4\pi\epsilon_0} \right)^2 \left[\ln \left(\frac{2m_e c^2 \beta^2}{I(1 - \beta^2)} \right) - \beta^2 \right], \quad (3.2)$$




FIGURE 3.7: Photo of a Silicon Tracking Telescope (STT), including the cooling system and read-out electronics.

where E is energy of the particle; x is the distance travelled by the particle; and m_e is mass of the electron; n is electron density of the target; I is the mean ionisation energy of the target; while ε_0 is the vacuum permittivity.

Therefore, these energy losses in the layers are specific to the particle type. Ratio of energy losses with the total energy of the particle is used to identify the particles. The minimum energy of a reconstructed particle is given by the thickness of the most inner layer. It will be detected as soon as it passes through the inner layer and in the second layer [40]. This corresponds to minimum energy of 2.5 MeV for protons. The setup was built in a ϕ -symmetric (left-right) arrangement to make use of the cross-ratio method. This configuration fulfils the requirement of particle identification together with a precise energy determination (1-5%) and tracking with vertex resolution of 1 mm over a wide range. The time resolution of the setup is less than 1 ns.

3.4 Experimental conditions

This thesis comprises data gathered over three dedicated experiments.

- Objective: analysing power $A_y(\theta)$ in proton-proton elastic scattering
 - Beam: polarised proton beam;
 - Target: unpolarised hydrogen cluster target;
 - Data gathered by: EDDA, ANKE STT, ANKE FD;

- Beam Energies: $T_p = 0.796, 1.6, 1.8, 1.965, 2.157, \text{ and } 2.368$ GeV;
- Angular range: 4° - 30° .
- Objective: unpolarised differential cross section in proton-proton elastic scattering $d\sigma/d\Omega$
 - Beam: unpolarised proton beam;
 - Target: unpolarised hydrogen cluster target;
 - Data gathered by: ANKE FD, Schottky spectrum analyser;
 - Beam Energies: $T_p = 1.0, 1.6, 1.8, 2.0, 2.2, 2.4, 2.6, \text{ and } 2.8$ GeV;
 - Angular range: 12° - 30° .
- Objective: analysing power $A_y(\theta)$ in proton-neutron quasi-free elastic scattering
 - Beam: polarised proton beam;
 - Target: unpolarised deuterium cluster target;
 - Data gathered by: EDDA, ANKE STT, ANKE FD;
 - Beam Energies: $T_p = 0.796, 1.6, 1.8, 1.965, 2.157, \text{ and } 2.368$ GeV;
 - Angular range: 13° - 30° .

Chapter 4

Analysing power in proton-proton elastic scattering

The investigations of the analysing power A_y in proton-proton ($\vec{p}p$) elastic scattering were motivated by the lack of experimental data at small angles above 1 GeV (Figure 1.3). The $\vec{p}p$ experiment at ANKE was carried out using polarised proton beam at six energies, $T_p = 0.796, 1.6, 1.8, 1.965, 2.157,$ and 2.368 GeV. The determination of the analysing power $A_y(\theta)$ as the function of the scattering polar angle requires the measurement of the scattered particles asymmetry $\varepsilon(\theta)$ and the beam polarisation value P :

$$\varepsilon(\theta) \propto PA_y(\theta). \quad (4.1)$$

While asymmetry $\varepsilon(\theta)$ was calculated using the ANKE detection systems (STT and FD), the beam polarisation P was measured using the EDDA detector. The beam polarisation mode was changed every subsequent cycle to take the maximum advantage of the cross-ratio method, described in Section 2.3. Cycles of 180 s or 300 s duration were used for each spin mode, with the last 20 s of each cycle being reserved for the measurement of the beam polarisation with the EDDA detector. The details on the beam polarisation measurement are provided in Section 4.1, while the asymmetry determination using the STT and FD are given in Section 4.2. Finally, the results of the analysing power in $\vec{p}p$ elastic scattering are discussed in Section 4.4.

4.1 Beam polarisation measurement using EDDA

The absolute values of the beam polarisations were measured by the EDDA polarimeter for the first time in the ANKE beam time in 2013. The EDDA carbon fibre target was moved into the beam for the last 20 seconds of every cycle, and scaler counts were recorded. The carbon target effectively consumes all the beam, hence it could not be used before the ANKE measurement of asymmetry in a cycle.

Originally, the EDDA detector was equipped with a polarised atomic hydrogen target, and had been used to measure the $p\vec{p}$ analysing power over almost the whole COSY energy range [12] [13]. By studying further the scattering of polarised protons on C and CH₂ targets, it was possible to deduce the quasi-free analysing power of the carbon.

The beam polarisation in our experiment has been determined based on the asymmetry $\varepsilon(\theta_{lab})$ and effective carbon analysing powers $A_n^{eff}(C)$ in quasi-elastic scattering of the protons on carbon-bound nuclear protons. These effective analysing power values were taken from the dedicated studies in 2000 at EDDA-COSY [34].

The asymmetry term $\varepsilon(\theta_{lab})$ was calculated via the cross-ratio method (Equation 2.19), using the rates in the semi-rings placed to the left and right of the beam. It was possible to compare left and right count rates for each polar angle θ_{lab} range, corresponding to the semi-ring (Appendix A), while averaging over azimuthal angle ϕ range. Thus, the asymmetry term for each polar angle range has form:

$$\langle \varepsilon(\theta_{lab}) \rangle = \langle PA_y(\theta_{lab}) \cos \phi \rangle \approx \frac{2}{\pi} PA_y(\theta_{lab}). \quad (4.2)$$

We can deduce the beam polarisation P from $\varepsilon(\theta_{lab})$ in an EDDA ring:

$$P_{\theta_{lab}} \approx \frac{\varepsilon(\theta_{lab})}{A_{eff}(\theta_{lab})}, \quad (4.3)$$

where A_{eff} is the effective analysing power, values for which are taken from [34]

$$A_{eff}(\theta_{lab}) = \frac{2}{\pi} A_y(\theta_{lab}). \quad (4.4)$$

The polarisation values calculated from the count asymmetry in every ring are shown in Figure 4.1 as the function of the corresponding scattering polar angle in the laboratory coordinate system. The beam polarisation P , obviously, does not

depend of the scattering angle θ_{lab} , at which we are performing the measurement. In order to decrease the statistical uncertainty of P we take advantage of this independence and form the weighted average of all $P_{\theta_{lab}}$ as a final result for P .




FIGURE 4.1: The polarisation values, calculated ring by ring, are shown versus corresponding laboratory polar angles (according to Appendix A). The sample plots are shown for beam kinetic energies $T_p = 1.8$ GeV and $T_p = 2.157$ GeV.

The weighted averages over time and polar angle of the beam polarisations determined at the six energies are given in Table 4.1. The changes in sign reflect the number of spin flips required to pass through the imperfection resonances, described in detail in Section 3.1.2. It should be noted that each of the six beams was prepared independently and, for this reason, the magnitude of the polarisation may not decrease monotonically as more resonances are crossed.

T_p (GeV)	0.796	1.6	1.8	1.965	2.157	2.368
P	0.554	0.504	-0.508	-0.429	-0.501	0.435
ΔP	± 0.008	± 0.003	± 0.011	± 0.008	± 0.010	± 0.015

TABLE 4.1: The values of the mean polarisations P determined with the EDDA polarimeter averaged over all the data at the beam energy T_p . Only statistical errors are given in the table.

Consistent results were achieved with the EDDA polarimeter after the short (180 s) and long (300 s) cycles. As expected, it implies that beam polarisation is not lost at flattop (constant beam energy) over a COSY cycle. However, due to the non-zero dispersion combined with the energy loss of the beam caused by its passage through the target, the settings at the three lowest energies gradually degrade slightly. This effect was taken into account in the analysis.

4.1.1 Beam polarisation uncertainty

The variation of the beam polarisation values among the cycles at any given energy was checked using EDDA with various selections and combinations of the cycles. All the studies yield the consistent results within the uncertainties at every given energy. The variation of the beam polarisation values was also checked with the asymmetry of the counts in the STT in various cycles and found to be around 0.04 (RMS).

Besides the statistical errors, shown in Table 4.1, the uncertainty of the effective carbon analysing powers should be taken into account. In the studies of the CH₂ and C targets for the fast beam polarisation, the systematic error for the A_{eff} , unfortunately, could be estimated only very roughly from the change of polarisation values during the calibration procedure. Unlike CH₂ target, C target is free of additional systematic errors occurring due to the aging of the target (loss of hydrogen ions in the interaction with beam protons). The value for the systematic uncertainty of the carbon effective analysing powers was estimated in [34] to be 3%.

4.2 Asymmetry measurement using ANKE

4.2.1 Event selection by STT

The analysis of the scattered particles, detected in the STT, is based on the software, developed mainly by G. Macharshvili. It includes the codes for reconstructing tracks, using energy deposited in the layers of the STT as well as the neural network method, and is described in detail in Reference [41].

Tracks were reconstructed starting from the hits in the second layer. The STT trigger requested a minimum energy deposit in the second layer of either telescope. Combinations of these hits with the hits in the first layer have been considered. If the reconstructed track hits the inside the ellipse of beam-target overlay in the zy plane (at $x = 0$), then it is stored.

In general, the hit from the third layer is also added to the reconstructed track in case it is inside the 20° cone along the track with the apex at the second layer hit. The cone opening angle corresponds to the maximum angle of multiple scattering.

The third layer hit does not change a track geometry, it is used only to fulfil the energy deposit information.

In some cases a possible track might be missing a hit in the first or second layer due to possible vicinity to the energy threshold or inactive segments. In order to increase the efficiency of the track reconstruction, in these cases energy deposits for the first or second layers were added to the energy loss sum in the correspondence to the missing hit, assuming $\Delta E_2 = 5\Delta E_1$ or $\Delta E_1 = 0.2\Delta E_2$ [42].

The greater precision in the angle of the recoiling proton is achieved by deducing it from the energy measured in the telescope rather than from a direct angular measurement. The Figure 4.2 demonstrates the difference between the angle reconstructed from the energy of the scattered particle and directly measured scattering angle of the track. Even though it is not possible to judge directly from Figure 4.2 the contribution of each angle to the distribution, the simulation shows that about 80% of the width of the distribution comes from the directly measured scattering angle of the track.




FIGURE 4.2: Difference between the directly measured angle of the track θ and the angle reconstructed from the energy $\theta(E)$. The example plot is shown at $T_p = 2.368$ GeV.

The protons with energies below 30 MeV were completely stopped in three layers of the STT. Furthermore, it is also possible to deduce the energy of punch-through protons up to 90 MeV. Thus expanding considerably the angular coverage of the telescope. For this purpose, the kinetic energy of these energetic protons was defined through a comparison of the angles and energy deposits in all three layers with simulated data using a neural network approach. The relative uncertainty

was defined during the network training procedure [43] and is equal to 2% at 30 MeV and 4% at 90 MeV.

There is very little ambiguity in the isolation of the proton peak in the missing mass spectra of the selected STT events (Figure 4.3).




FIGURE 4.3: Missing mass $M_X(pp \rightarrow pX)$ spectrum obtained for the beam energy of $T_p = 1.6$ GeV, showing the clear proton peak when detecting one proton in the STT.

4.2.2 Asymmetry measurement using STT

The left-right symmetry of ANKE STTs, along with the reversal of the beam polarisation every subsequent cycle allowed us to use the cross-ratio method, described in the Section 2.3. The left-right asymmetry of the protons scattered to the STTs in each polar angle interval is calculated as

$$\varepsilon(\theta) = \frac{L(\theta) - R(\theta)}{L(\theta) + R(\theta)}, \quad (4.5)$$

where $L(\theta)$ and $R(\theta)$ are the geometrical means of number of particles scattered in the given polar angle interval to the left $L = \sqrt{N_1^\uparrow N_2^\downarrow}$ and to the right $R = \sqrt{N_1^\downarrow N_2^\uparrow}$ in respect to the beam polarisation (Figure 2.2). As was shown previously, the scattering asymmetry is related to the analysing power $A_y(\theta)$ for each value of the scattering angle θ through

$$\varepsilon(\theta) = A_y(\theta)P\langle \cos\phi \rangle. \quad (4.6)$$

The ϕ acceptances for the left and right STTs differ only slightly: $\langle \cos\phi \rangle_1 = 0.9663 \pm 0.0005$ and $\langle \cos\phi \rangle_2 = 0.9670 \pm 0.0003$. The simulation showed that even larger difference between $\langle \cos\phi \rangle_1$ and $\langle \cos\phi \rangle_2$ would not affect the measured asymmetry. Therefore, we assume that ϕ acceptances of the detectors are the same and in our calculations $\langle \cos\phi \rangle \approx 0.9666$ can be used.

4.2.3 Event selection by FD

Scattered particles, which pass through the vacuum chamber of the D2 magnet and leave it through the forward exit window, are detected in the FD. The analysis is based on the software, developed mainly by S. Dymov. It includes the codes for finding tracks and reconstructing particle momenta and is described in detail in Reference [37].

The precision of the momentum and angle reconstruction is directly related to the accuracy of the ANKE geometry measurement. Positions and sizes of various parts of the ANKE spectrometer are well defined and fixed, but there are some parameters in the track reconstruction software that change from one experiment to another and it is not possible to measure them directly with enough precision. Uncertainties in these parameters shift the reconstructed particle momenta and consequently result in shifts in the missing mass spectra. Therefore, these parameters should be calibrated for every experiment individually.

The setup parameters are adjusted using the kinematics of different reactions. For every iteration of the fitted parameters, the program reconstructs tracks from scratch and looks at the displacement of the missing masses from their nominal values for every reaction. After the geometry adjustment, the hit coordinates and the time of flight are used for the final track reconstruction. The energy loss cut for elastic proton identification is relevant only at the 0.796 GeV beam energy, where the forward going deuterons from the $pp \rightarrow d\pi^+$ have the momenta close to that of the pp elastic protons. After the applied cuts the admixture of those deuterons does not exceed 0.02%.

The number of elastic protons was determined from the missing mass spectra after subtracting a linear background from the peak in each angular bin. Typical missing mass spectrum is shown in Figure 4.4.

The setup acceptance was defined from GEANT simulation, followed by the same track and momentum reconstruction procedures that were applied to the data.




FIGURE 4.4: Missing mass $M_X(pp \rightarrow pX)$ spectrum obtained from the particles detected in the forward detector at the beam kinetic energy of 1.6 GeV.

Although the final results were obtained without further restriction on the ϕ range, estimations done with several ϕ cuts showed no change beyond the statistical fluctuations. The angular acceptance at $T_p = 1.6$ GeV is shown in Fig. 4.5 and it looks rather similar at the other energies [44].




FIGURE 4.5: Forward detector angular acceptance for $pp \rightarrow pp$ at the beam kinetic energy of 1.6 GeV.

4.2.4 Asymmetry measurement using FD

The absence of the left-right symmetry in the Forward Detector does not permit the use of the cross-ratio method to determine the asymmetry. Therefore, the

analysing power can only be defined from the asymmetry of the count rates for the two states of the beam polarisation. In this case the asymmetry is introduced as

$$\varepsilon(\theta) = \frac{N_{\uparrow}(\theta)/L_{\uparrow} - N_{\downarrow}(\theta)/L_{\downarrow}}{N_{\uparrow}(\theta)/L_{\uparrow} + N_{\downarrow}(\theta)/L_{\downarrow}} \quad (4.7)$$

in terms of the luminosity-normalised numbers of counts for the two orientations of the beam polarisation $N_{\uparrow}(\theta)/L_{\uparrow}$ and $N_{\downarrow}(\theta)/L_{\downarrow}$, or

$$\varepsilon(\theta) = \frac{N_{\uparrow}(\theta) - N_{\downarrow}(\theta)/L_{rel}}{N_{\uparrow}(\theta) + N_{\downarrow}(\theta)/L_{rel}} \quad (4.8)$$

in terms of the relative luminosity factors $L_{rel} = L_{\uparrow}/L_{\downarrow}$. These factors were fixed by comparing the rates of charged particle production in angular regions where the beam polarisation could play no part [45].

The luminosity calibration data, which corresponded generally to the inelastic events involving pion production, were selected by applying cuts either on the small polar angles θ or on the azimuthal angle ϕ near $\pm 90^\circ$. In Figure. 4.6 three groups of events with the decreasing level of cut tightness are shown:

- i) $\theta < 0.5^\circ$, $||\phi| - 90| < 5^\circ$
- ii) $\theta < 1^\circ$, $||\phi| - 90| < 10^\circ$
- iii) $\theta < 2^\circ$, $||\phi| - 90| < 20^\circ$.




FIGURE 4.6: Relative luminosities obtained with various angular cuts (shown at horizontal axis) at $T_p = 1.8$ GeV. Only a small section of the vertical scale is shown to emphasise the small differences.

Figure. 4.6 illustrates the relative luminosities at the sample energy $T_p = 1.8$ GeV, but the distributions look similar at the other energies. Luminosity ratios in the three group of cuts are consistent within the statistical uncertainty. For the calculation of the asymmetry, the average value of the relative luminosity factors, gained from the abovementioned cuts, was used as the normalisation factor in Equation 4.8.

4.3 Sources of the systematic uncertainties

The cross-ratio method allows one to eliminate first order systematic errors that arise from misalignments between the left and right STTs. Other systematic errors for the asymmetry obtained using the STT, such as those arising from differences in the magnitudes of the up and down polarisations, also cancel in the first order. Nevertheless, the systematic uncertainties of the STT data were investigated further to the higher order by varying all the inputs to the reasonable levels, and observing the possible change in the asymmetry. The data gained using the FD lack these advantages and rely on the correct normalisation with the relative luminosities. The three groups of angular cuts, described previously, yielded consistent results within statistical errors. As a result, it is estimated the systematic uncertainty of A_y due to the normalisation procedure does not exceed 0.3%.

4.3.1 Difference in polarisation values for two polarisation modes

Low Energy Polarimeter (LEP) measurements showed that at the injection into the COSY ring, the polarisation magnitudes for polarisation modes “up” and “down” are less than statistical uncertainty of 1%. After the acceleration the difference between the values for two modes may be larger. However, even assuming $|P_{\uparrow} - P_{\downarrow}| = 0.1$, the correction term for the asymmetry measured using the STT is less than $5 \cdot 10^{-4}$ and can be ignored. The difference between the polarisation magnitudes is potentially more serious for the FD analysis, the same assumption of $|P_{\uparrow} - P_{\downarrow}| = 0.1$ could induce the relative errors in A_y of up to 2.5%. However, it should be noted that in the overlap regions of the STT and FD data any disagreements between the determinations of the asymmetries in the two systems are on the 1% level and this puts a much tighter constraint on possible $|P_{\uparrow} - P_{\downarrow}|$ differences.

4.3.2 Polar angle accuracy

Misalignment of the left and right STT detectors, *i.e.* the difference in the angles θ_1 and θ_2 measured at two telescopes, may raise the fake asymmetry. Assuming $|\theta_1 - \theta_2| \simeq (0.5 \pm 1.0)^\circ$ for each θ bin, estimated from $\theta - \theta(E)$ distribution variance (Figure 4.2), the upper limit for the systematic error induced by the STT detectors misalignment is $1.5 \cdot 10^{-3}$ at the beam kinetic energy of $T_p = 0.796$ GeV and by factor of 2 smaller at the higher energies.

One can estimate the precision of the measured scattering angle at the FD indirectly from the $pp \rightarrow d\pi^+$ reaction. The simultaneous measurement of the deuteron and pion in the FD showed that the precision of the horizontal transverse momentum ΔP_x is less than 1.5 MeV/c. This corresponds to the systematic deviation in the laboratory angles from those expected for these kinematics $\Delta\theta_{lab} < 0.07^\circ$. If this is valid also for pp elastic scattering it would suggest that the c.m. scattering angles were defined with a precision of better than 0.15° .

In cases, where one of the protons from an elastic scattering event is detected in the FD and the other in the STT, it is possible to compare directly the scattering angle measured in the two systems. Figure 4.7 shows the difference of the results of the θ_{cm}^{STT} calculated from the energy deposited in the STT layers and θ_{cm}^{FD} reconstructed from the track in the FD.

In general, the θ_{cm}^{STT} angle is larger than θ_{cm}^{FD} , with the difference being typically $\approx 0.3^\circ$, as seen for $T_p = 0.796$ GeV (left panel) and $T_p = 2.157$ GeV (right panel). It is not possible to judge which detector is responsible for this difference which is, however, small compared with the θ_{cm} bin widths of 1.0° (FD) and 1.2° (STT) used for the final values of A_y .




FIGURE 4.7: Difference between the scattering angles reconstructed using the FD θ_{cm}^{FD} and STT θ_{cm}^{STT} . Example plots are shown for $T_p = 0.796$ GeV (left panel) and $T_p = 2.157$ GeV (right panel).

To estimate the impact of the angular deviation onto A_y , for each energy we approximated the reconstructed A_y dependence with a parabola. Then, one can calculate the relative change of A_y occurring due to the constant shift in angle $\delta A_y = (A_y(\theta) - A_y(\theta + \delta\theta))/A_y(\theta)$ (Figure 4.8). The maximal error of 2.5% is obtained at $T_p = 0.796$ GeV. At higher energies the possible systematic error is much smaller, *e.g.*, at $T_p = 2.368$ GeV, shown in the right panel of Figure 4.8, it is less than 1.2%.




FIGURE 4.8: Systematic error due to the possible maximum shift of the scattering angle. Example plots are shown for the beam kinetic energies $T_p = 0.796$ GeV (left panel) and $T_p = 2.368$ GeV (right panel).

4.3.3 Detector efficiencies stability

Another factor that could affect the asymmetry measured using the STTs is an instability of two detectors efficiencies in time. To be more precise, the ratio of the efficiencies of the two telescopes $r = E_1/E_2$ should be constant to avoid the fake asymmetry. Let us introduce the instability factor $r_{\uparrow}/r_{\downarrow}$ that should be equal to 1, in case the efficiencies ratios are the same for the both beam polarisation modes.

The angular dependence of typical average ratio $r_{\uparrow}/r_{\downarrow}$ at the beam energies of $T_p = 0.796$ GeV and 1.8 GeV are shown in Figure 4.9. In almost all the cases the fitted constants are indeed close to 1 (with $\chi^2/ndf \simeq 1$).

The uncertainty of the fitted constant can be interpreted as the systematic uncertainty induced by the detector instability, *i.e.*

$$\Delta A_y(\theta) \approx (1 - r_{\uparrow}/r_{\downarrow})A_y(\theta) \equiv c(\theta)A_y(\theta). \quad (4.9)$$




FIGURE 4.9: The angular dependences of the instability factor $r_{\uparrow}/r_{\downarrow}$ at the beam energies of $T_p = 0.796$ GeV (left panel) and $T_p = 1.8$ GeV (right panel).

The STT detector efficiency instability correction, which was studied at all energies, does not exceed the $|c| = 1.3\%$ that was found at 1.8 GeV. The relevant corrections of the analysing power $c(\theta)A_y(\theta)$ were added for each angular bin of the analysing power, obtained from the STT data set.

In order to simulate the maximum possible systematic error, induced by the STT detectors inefficiencies, $c = 1 - r_{\uparrow}/r_{\downarrow} = 0.2$ was introduced into the simulation. After the correction factor was applied, the angular dependence of the systematic uncertainty has been simulated once more, and the obtained systematic uncertainties were typically of the order 10^{-3} that is one order smaller than prior to the correction.

The efficiency for registering events in the FD were studied by using events where both the fast and recoil protons were measured in the FD and STT, respectively. The efficiencies obtained per cycle are shown in Figure 4.10, summarised separately for the two beam polarisation modes. This value is expected to be lower than 100% due to a small fraction of misidentified protons in the STT, as well as the possibility of multiple particles recorded in the STT or FD within a single trigger. There is also a small inefficiency of the FD trigger ($\lesssim 1\%$) and a loss of particles because of the large angle scattering or interaction with the detector material.

Most importantly, the difference of the efficiencies between the two polarisation modes is of less than 10^{-3} that is within the statistical uncertainties.




FIGURE 4.10: Efficiencies of the elastic event reconstruction using the FD in the cycles with the beam polarisation mode up (left panel) and beam polarisation mode down (right panel).

4.3.3.1 Summary on systematic uncertainties

The overall systematic uncertainty in A_y arising from asymmetry measurement using the STT does not exceed 0.3%. The dominant systematic error is, hence, that arising from the determination of the beam polarisation in the EDDA polarimeter, which was estimated to be 3% [34]. For the FD data there is, in addition, a possible contribution associated with the assumption of equal up and down polarisations so that in this case we would cautiously assume a 5% systematic uncertainty. To these figures must be added the statistical uncertainty in the determinations of the beam polarisations at the six energies shown in Table 4.1.

4.4 Results and discussion

The results of all the ANKE measurements of $A_y(\theta_{cm})$ for $\vec{p}p$ elastic scattering are shown for the six energies in Figure 4.11. For the final results, only those cycles were used that contained good quality data in all the detectors: FD and STT, as well as the EDDA polarimeter.

The agreement between the STT and FD data, which involved completely independent measurements of the final state, is remarkably good. The individual deviations generally lie within the statistical uncertainties and the average Over the angular overlap regions is $A_y(FD)/A_y(STT) = 1.00 \pm 0.01$. At beam energies close to 0.796 GeV there are many measurements of the pp analysing power and, in general, they are in good mutual agreement, as they are with the new ANKE data.




FIGURE 4.11: Comparison of the ANKE measurements of the proton analysing power in $\vec{p}p$ elastic scattering using the STT (red filled circles) and the FD (blue filled triangles) systems with the curves corresponding to the SAID 2007 (solid black line) [8] and the revised fit (dashed red) solutions. Only statistical errors are shown so that the systematic uncertainties arising, for example, from the calibration of the EDDA polarimeter have not been included. Also shown are selected results from EDDA (black crosses) [13] at the energies different by no more than 7 MeV and, at 0.796 GeV, LAMPF [46–48], and SATURNE [49] (black open symbols).

This reinforces the confidence in the use of the EDDA polarimeter. At 1.6 GeV and above there are far fewer experimental measurements and, for clarity, we only show the EDDA data at neighbouring energies though, at the highest energy, the statistical fluctuations are significant [13].

The SAID 2007 solution [8], shown by the solid black line in Figure 4.11, describes the bulk of the ≈ 0.796 GeV data very well indeed. However, at higher energies the ANKE data deviate significantly from the predictions of the 2007 solution. Moreover, the shapes of the ANKE data seem very different from these predictions, rising much more steeply at small angles. Therefore, these discrepancies cannot be due to a simple miscalibration of the EDDA polarimeter, for example, which would change the overall magnitude of $A_y(\theta)$ but not its angular dependence [50].

The ANKE analysing power data have been added to the world data set and searches made for an updated partial wave solution. To allow for possible systematic effects, the SAID fitting procedure introduces a scale factor N into any

data set and determines its value, as well as the pp phases and inelasticities, by minimising an overall χ^2 for the complete data set. When this is done, the average value of χ^2 per degree of freedom found for the ANKE STT data is 1.6 and slightly larger for the FD results. The new fits, which lead to the red dashed curves in Figure 4.11, correspond to significant changes at the higher energies to the parameters of the lower partial waves, with the biggest changes up to 60% in 3F_2 and 3F_4 (Figure 4.12).




FIGURE 4.12: Energy-dependent phase shift analysis parameters: δ phase shift and ρ ($\cos \rho = \eta$, where η is the absorption parameter) for 3F_2 and 3F_4 partial waves from the 2014 SAID solution, in comparison with the older solution from 2007.

The values of the normalisation factors N reported in Table 4.2 have an average of $\langle N \rangle = 1.00 \pm 0.02$ for the STT data. These factors, which would effectively multiply the beam polarisations, have not been applied in Figure 4.11. The deviations of the individual values of N from unity might seem to be greater at the higher energies. They are somewhat larger than what one would expect on the basis of the quoted uncertainties in the EDDA polarimeter, being around 5% rather than the 3% estimate [34]. It should be stressed that the introduction of the scale factor N does not change the shape of a distribution and, even in cases where a value close to one is found, this does not mean that the fit reproduces perfectly the data. A clear example of this is to be found in the larger angle data at 1.6 GeV shown in Figure 4.11.

T_p (GeV)	0.796	1.6	1.8	1.965	2.157	2.368
N	1.00	1.00	0.99	1.09	1.01	0.93

TABLE 4.2: The normalisation factors N obtained in a partial wave fit [51] to the current STT data.

4.5 Conclusion

We have measured the analysing power in $\vec{p}p$ elastic scattering at 0.796 GeV and at five energies from 1.6 GeV up to 2.4 GeV using both the silicon tracking telescopes and the forward detector. The consistency between these two independent measurements of the final protons is striking so that the only major systematic uncertainty is the few percent associated with the calibration of the EDDA polarimeter. Though the overall uncertainties are slightly larger for the FD data, these results are important because they extend the coverage to slightly larger scattering angles.

In the small angle range accessible to ANKE, the new data are consistent with older measurements around 0.796 GeV and also with the 2007 SAID predictions at this energy [9]. At higher energies the ANKE results lie significantly above the 2007 solution near the forward direction and also display a different angular dependence. By adjusting some of the phases and inelasticities in the low partial waves by up to 60%, the new SAID solution [51] was obtained that provides a much better description of the new ANKE A_y data.

Chapter 5

Cross section in proton-proton elastic scattering

As was shown in the introduction (Figure 1.3), there are relatively few measurements of pp elastic scattering for beam energies above about 1 GeV in the angular range between 10° and 30° . This falls between the region of major Coulomb effects and the larger angles where the EDDA collaboration has contributed extensively. This lack of information on the differential cross section and the analysing power inevitably leads to ambiguities in any pp PWA at high energies. To address this gap in our knowledge, ANKE collaboration has carried out the studies of the pp analysing power, presented in the previous chapter, and differential cross section, which is discussed here.

The differential cross section $d\sigma/d\Omega$ measurements were performed using the ANKE unpolarised hydrogen cluster-jet target and the unpolarised proton beam at eight kinetic energies $T_p = 1.0, 1.6, 1.8, 2.0, 2.2, 2.4, 2.6, 2.8$ GeV for center-of-mass angles in the range from 12° - 16° to 25° - 30° , depending on the energy. As was the case for the analysing power A_y , the present studies of $d\sigma/d\Omega$ were carried out using the ANKE spectrometer. However, for $d\sigma/d\Omega$ investigations only forward detection system was used to detect fast protons from elastic pp scattering.

In general, the cross section σ for a given physical process is given in terms of the corresponding counting rate R and the luminosity L through:

$$\sigma = \frac{R}{L} . \tag{5.1}$$

The biggest challenge is to establish the beam-target luminosity at the few percent level, because the overlap of the beam and target can not be deduced with such precision from macroscopic measurements. The details on the luminosity determination are provided in Section 5.1, while the count rate analysis using FD is shortly described in Section 5.2. Finally, the results of the differential cross section in pp elastic scattering are discussed in Section 5.4.

5.1 Luminosity determination

The knowledge of the luminosity is the crucial element of cross section measurement. The ANKE collaboration and the COSY machine crew have jointly developed a very accurate method for determining the absolute luminosity in an experiment at an internal target position. The technique relies on measuring the energy losses due to the electromagnetic interactions of the beam as it passes repeatedly through the thin target and measuring the shift of the revolution frequency by studying the Schottky spectrum [52]. In order to extract the cross section of any reaction, the absolute value of the luminosity must be determined. For the fixed target experiments, luminosity is completely defined by:

$$L = n_B \times n_T, \quad (5.2)$$

where n_B is the particle current of the incident beam and n_T is the effective target thickness, expressed as an areal density. The measurement of the beam intensity n_B is performed via the high precision Beam Current Transformer (BCT) (Section 5.1.1). The effective target density n_T is determined via the measurement of the beam revolution frequency shift caused by the electromagnetic interaction between the beam and target particles (Section 5.1.2).

5.1.1 Beam intensity measurement

The particle current of the incident beam n_B was deduced from the calibrated Beam Current Transformer signal. The amplitude of the BCT signal is proportional to the COSY beam current, $I_b = n_B e = N f e$, where N is number of particles in the beam, f is revolution frequency, and e is particle charge.

To avoid the effects from stray magnetic fields, the BCT is mounted in field-free region of the ring and, in addition, is magnetically shielded.




FIGURE 5.1: Beam Current Transformer (BCT) typical raw signal, recorded during the experiment.

The raw BCT signal (Figure 5.1) was calibrated with a current-carrying wire placed between the beam tube and ferrite core of the BCT. The result of the calibration, applying a current from a high precision source, is given in Figure 5.2 and it does not depend on the energy of the beam.




FIGURE 5.2: BCT signal vs. current I in the calibration wire.

In the given experiment, the BCT signal was recorded directly to the ANKE data stream. The measurements were carried out every second over the 300 s cycle and then averaged. The final results are accurate to better than 10^{-3} , and hence, can be neglected in comparison with target density uncertainties that will be discussed in the following.

5.1.2 Target density determination

When a charged particle passes through the matter, it loses energy through electromagnetic interaction. In the storage ring an uncooled beam passes a thin target a very large number of times. The energy loss, which is proportional to the target thickness, builds up steadily in time and causes a shift in the revolution frequency f_0 . This shift can be determined by studying the Schottky power spectrum of the beam [53].

The energy loss δT per single passage through the target, divided by the mass stopping power $S_m = \frac{dE}{dx} \frac{1}{\rho}$ and the mass m of the target atom, yields the number of target atoms per unit area that interact with the beam:

$$n_T = \frac{\delta T}{S_m m}. \quad (5.3)$$

Over a small period of time the beam makes $f_0 \Delta t$ traversals with the corresponding energy loss ΔT . Hence, Equation 5.3 maybe written in terms of energy loss, as

$$n_T = \frac{\Delta T}{f_0 S_m m \Delta t}, \quad (5.4)$$

or applying logarithmic differentiation, in terms of change in the beam momentum as

$$n_T = \frac{1 + \gamma}{\gamma} \frac{T_0 \Delta p}{f_0 S_m m p_0 \Delta t}. \quad (5.5)$$

T_0 and p_0 are the initial values of the beam kinetic energy and momentum and γ is the Lorentz factor.

The fractional change in the revolution frequency is proportional to that in the momentum through the so-called frequency-slip parameter η :

$$\frac{\Delta f}{f_0} = \eta \frac{\Delta p}{p_0}. \quad (5.6)$$

Putting the expressions 5.5 and 5.6 together, we obtain

$$n_T = \left(\frac{1 + \gamma}{\gamma} \right) \frac{1}{\eta} \frac{1}{S_m m} \frac{T_0}{f_0^2} \frac{df}{dt}. \quad (5.7)$$

The stopping power S_m and mass of the target atoms are well known and easily accessible at NIST-PML data base [54]. The initial values of frequency and energy of the beam, as well as the Lorenz factor are routinely measured for every

experiment at COSY. Therefore, once the frequency-slip parameter η is known, n_T can be deduced by measuring the beam revolution frequency shift df/dt during the data taking.

The frequency-slip parameter η can be expressed in terms of α , the so-called momentum compaction factor, which is constant for a given lattice settings of the accelerator:

$$\eta = \frac{1}{\gamma^2} - \alpha. \quad (5.8)$$

The η parameter determines the direction of the frequency change during the machine cycle. The revolution frequency depends on the particle speed βc and the orbit length L through $f = \beta c/L$, where due to dispersion, L is also a function of the momentum. Using the definition of the momentum compaction factor $\alpha = (dL/L)/(dp/p)$, we obtain

$$\frac{df}{f} = \left(\frac{1}{\gamma^2} - \alpha \right) \frac{dp}{p}. \quad (5.9)$$

An estimate of α , which is constant for a given lattice setting, may be done by a computer simulation of the accelerator, but greater precision is achieved by a direct measurement, as will be described later.

5.1.2.1 Schottky noise

The beam in the synchrotron consists of a finite number of charged particles. A current created by these charge carriers has some statistical fluctuation, that were investigated first by W. Schottky [55].

The current fluctuations induce a voltage signal at a beam pick-up. The Fourier analysis of this voltage signal by a spectrum analyser delivers frequency distribution around the harmonics of the revolution frequency. The frequencies of COSY were measured with the existing Schottky pick-up of stochastic cooling system, which is optimized to operate in GHz region. The harmonic number 1000 of COSY revolution frequency was measured with a spectrum analyser. Schottky noise current is proportional to the square root of number of particles in the ring. Therefore, the amplitudes of the measured distribution are squared to give the Schottky power spectra, which are representative of the momentum spread of the beam.




FIGURE 5.3: Schottky power spectra for $T_p = 1.0$ GeV (a) and $T_p = 2.0$ GeV (b) obtained during one 300 s cycle and scaled to harmonic number 1. The mean frequencies are indicated by the vertical (red) lines.

The Schottky signals were recorded every 10 s with a 189 ms sweep time, resulting in effectively instantaneous spectra. For ease of representation, only the results from every 60 seconds are shown in Figure 5.3.

We estimate the background from both sides of the spectra, assuming that the background noise associated with a spectrum analyser is flat. After subtracting the background noise from the original Schottky spectrum, we can evaluate the mean frequency f of the beam at any instantaneous measurement. The centroids of the distributions, are indicated by the vertical red lines in Figure 5.3.

5.1.2.2 Frequency shift

The mean frequencies of the beam as the function of time t over the 300 s cycle are presented in Figure 5.4. It must be noted that the direction of the frequency change is actually different at low and high energies, depending on the lattice settings of the accelerator. The point of transition, where df changes its sign,




FIGURE 5.4: Typical mean frequency shift derived from the Schottky power spectra of the type illustrated in Figure 5.3 for $T_p = 1.0$ GeV (left panel) and $T_p = 2.0$ GeV (right panel).

occurs when $\alpha = 1/\gamma_{tr}^2$. Consequently, near the transition point η is small and this is the principal restriction on the applicability of the Schottky method. For COSY proton beam the transition occurs for $T_p \approx 1.3$ GeV, for this reason the experiments were not conducted between 1.0 GeV and 1.6 GeV. As one can see, an average frequency shift in the cycle, which is comparable to the spectra width, is negative for 1.0 GeV and positive for 2.0 GeV. For this reason, the example energies in Figures 5.3-5.5 were chosen from both sides of the transition point.




FIGURE 5.5: Average frequency shifts within the cycle duration, measured by the new (blue) and old (green) spectrum analysers at $T_p = 1.0$ and $T_p = 2.0$ GeV.

The frequency change df/dt should not be time dependent, if beam-target overlap and the target density are constant during the cycle. Hence, by checking df/dt dependence on time within the cycle via averaging many cycles, one can check the stability of the effective target density, which also accounts for beam-target overlap.

In order to avoid the systematic effects, associated with the COSY magnets slowly reaching the nominal field, it was decided to cut off the data from first 60 seconds

of data taking in every cycle. The frequency shifts were calculated independently, splitting the data from the useful cycle length into three parts. As demonstrated in Figure 5.5, df/dt values did not show any evidence of time dependency apart from the statistical fluctuations.

5.1.2.3 COSY rest gas

A small frequency shift is produced by the interaction of the beam with the residual gas in the COSY ring. Besides, the ANKE target produces additional increase of the vacuum pressure in the vicinity of the target. In order to account for these systematic effects, dedicated cycles were developed, where the ANKE cluster target was switched on, but the beam was steered away from it.

Even though these measurements already account for both effects, the separate cycles with target switched off, were performed, to estimate the pure effect of the residual gas, and to check that the beam was successfully steered away from the target. Figure 5.6 shows an example of these measurements at 1 GeV. As one can see, the effects from the ANKE target and residual gas together, as well as residual gas only are small, but distinguishable.




FIGURE 5.6: Example of frequency shift caused by the background when the target is on $\left(\frac{df}{dt}\right)_{ANKE+COSY}$ (left) and when the target is off $\left(\frac{df}{dt}\right)_{COSY}$ (right) compared to the total frequency shift $\left(\frac{df}{dt}\right)_{tot}$ with the beam incident on the target (middle).

Finally, the frequency shift caused purely by beam-target interaction is calculated as:

$$\left(\frac{df}{dt}\right)_{tar} = \left(\frac{df}{dt}\right)_{tot} - \left(\frac{df}{dt}\right)_{ANKE+COSY} \quad (5.10)$$

where $\left(\frac{df}{dt}\right)_{tot}$ is the total frequency shift during the measurement discussed in the previous section, and $\left(\frac{df}{dt}\right)_{ANKE+COSY}$ is the frequency shift caused by the residual gas of the COSY ring and effect from the ANKE cluster target. The corresponding uncertainties of the background measurement have been accounted for in the systematic errors summary (see details in Table 5.1).

5.1.2.4 Momentum compaction factor

After determining df/dt , only momentum compaction factor α is needed to finalise the calculation of the target density n_t at all the beam energies, according to Equation 5.7 and 5.8. The α measurements at COSY were performed in the dedicated cycles with the target switched off by adjusting the strength of the COSY bending magnets by few parts per thousand (which lays within the accelerator acceptance) and measuring the relative change in the beam revolution frequency $\Delta f/f_0$. The dependence between $\Delta B/B_0$ and $\Delta f/f_0$ is plotted in Figure 5.7.




FIGURE 5.7: Variation of the relative change in the mean beam revolution frequency $\Delta f/f_0$ with the relative change in the field strength in the bending magnets $\Delta B/B_0$. The fit defines the momentum compaction factor α . The example is shown for $T_p = 2.0$ GeV.

If the changes are not very large there is a linear relationship between the relative revolution frequency shift $\Delta f/f_0$ and the relative change in the field $\Delta B/B_0$,

expressed as following:

$$\frac{\Delta f}{f_0} = \alpha \frac{\Delta B}{B_0}. \quad (5.11)$$

Larger values of $\Delta B/B_0$ and $\Delta f/f_0$ require higher order terms in Equation 5.11. For the purposes of this experiment, $|\Delta B/B_0| < 1\%$ range was fitted with polynomial of second order.

5.1.2.5 Luminosity summary

The luminosity measurements defined by the effective target density n_T and beam current n_B are summarised for all eight energies of the experiment in Figure 5.8.




FIGURE 5.8: The luminosity measured in every cycle during the experiment.

Every point stands for the average luminosity in the cycle. As seen, the luminosity may vary from cycle to cycle rather significantly even at the fixed beam energy, which is mainly due to the COSY beam intensity variations. This does not concern the reliability of the final results, because the target density is determined for each cycle independently.

5.2 Event selection by FD

In order to determine cross section, besides luminosity counting rate R (Equation 5.2) of pp elastic scattering events must be measured. The events were selected by detecting a single fast proton in the Forward Detector (FD). The analysis of pp elastic scattering data from the FD has been already discussed in the previous chapter. Therefore, in this section only the specifics of the cross section analysis will be discussed.

The trigger for the data acquisition system was initiated by a signal produced by the proton in either of two hodoscopes of FD, placed one behind the other. This, together with a high efficiency of scintillation counter, reduced the trigger inefficiency to the 10^{-4} .

The counter efficiency was studied by analysing the signal amplitude as a function of the vertical hit coordinate for the selected pp elastic events. Two kinds of angular acceptance cuts were applied: the equal $|\phi| < 10^\circ$ cut for all polar angles, and $|Y_{track}| < 10$ cm cut on the vertical coordinate in the D2 exit window. No sign of the amplitude falling below the threshold was observed, except for the first (closest to the beam pipe) counters in each hodoscope wall. The efficiency of each of MWPC's planes exceeds 97% [56].

The selected events produced a prominent peak in the missing mass spectrum, with a background of only 1-2%. A small contribution from the $pp \rightarrow d\pi^+$ reaction to the peak region at 1 GeV was subtracted on the basis of the energy-loss information.

The setup parameters were adjusted in a geometry tuning procedure, with the use of the exclusive $pp \rightarrow pp$, $pp \rightarrow pn\pi^+$, $pp \rightarrow pp\pi^0$, and $pp \rightarrow d\pi^+$ reactions. In the last case, both the d and π^+ were detected in the FD simultaneously and this gave valuable information on the systematic uncertainty of the transverse momentum reconstruction. These showed that any systematic shift in the determination of the c.m. angle in pp elastic scattering was less than 0.15° , which would correspond to a 0.5% change in the differential cross section.

results are shown in Figure 5.9. Also shown are exponential fits to the measured data made on the basis of

$$\frac{d\sigma}{dt} = A \exp(-B|t| + C|t|^2), \quad (5.12)$$

where the values of the resulting parameters are given in Table 5.2.




FIGURE 5.9: Combined ANKE data set of differential cross sections with respect to the four-momentum transfer t compared to fits made on the basis of Equation 5.12. Systematic errors are not shown. The correct values are shown at 1.0 GeV but, for clarity of presentation, the other data are scaled down sequentially in energy by factors of 1.2. The true numerical values of the cross section and fit parameters are given in Appendix C and Table 5.2, accordingly.

Taking $C = 0$ at 1 GeV would change the value extracted from the fit by less than 1%, though this parameter becomes more important at higher energies where the t range is larger. This empirical representation of the measured data may prove helpful when the results are used in the normalisation of other experimental measurements.

5.4.1 Comparison with LNPI results at 1 GeV

There are very few data sets of absolute cross sections at small angles to which the ANKE results can be compared. In the vicinity of 1 GeV there are two measurements that were made with the IKAR recoil detector in the Leningrad Nuclear Physics Institute (LNPI). In the first of these at 992 MeV, IKAR used a hydrogen target [58]. In the second at 991 MeV a methane target was used, though the prime purpose of this experiment was to show that such a target gave consistent results and so could be used with a neutron beam [59].




FIGURE 5.10: Differential cross section for pp elastic scattering. The ANKE data at $T_p = 1$ GeV with statistical errors (blue squares) are compared to the IKAR (LNPI) hydrogen data at 992 MeV (green circles) [58] scaled by a factor of 1.085 and methane results at 991 MeV (red triangles) [60] scaled by a factor of 1.04. At very small values of $|t|$ there is a rise caused mainly by Coulomb-nuclear interference.

The ratio of the IKAR hydrogen values [58] to the fit of the ANKE 1 GeV data over the common range of angles is 0.920 ± 0.005 . In order to compare the shapes of these data sets, the LNPI results have been scaled by a factor of 1.085 before being plotted in Figure 5.10. The scaling factor is significant in view of the 2% and 2.8% absolute normalisations reported for the IKAR and ANKE experiments, respectively.

5.4.2 Comparison with ANL results at 2.0 and 2.8 GeV

Data are also available from the Argonne National Laboratory (ANL) in our angular range at 2.2 and 2.83 GeV [61] and these values are plotted together with our

measurements in Figure 5.11. The ANL data sets agree with our 2.2 and 2.8 GeV results to within 1%. However, the absolute normalisation claimed for these data was 4% [61] so that it is not possible to draw completely firm conclusions from this comparison.




FIGURE 5.11: The ANKE pp differential cross section data at 2.2 GeV (closed blue circles) and 2.8 GeV (closed blue triangles) compared to the ANL results [61] at 2.2 GeV (open red circles) and 2.83 GeV (open red triangles). Systematic errors are not shown. For presentational purposes, both higher energy data sets have been scaled downwards by a common factor of 1.5.

5.4.3 Impact on the SAID solution

The results obtained at ANKE could clearly have an impact on the current partial wave solutions. This is demonstrated in Figure 5.12, where the ANKE cross sections at 1.0, 2.0, and 2.8 GeV are compared to both the SAID 2007 solution [9] and a modified one [51] that takes the present data at all eight energies into account. Scaling factors in the partial wave analysis, consistent with the overall uncertainties given in Table 5.1, have been included in the figure. The major changes introduced by the new partial wave solution are in the 1S_0 and 1D_2 waves.

The precise EDDA measurements were undertaken for c.m. angles of 35° and above whereas the ANKE data finish well below this and the gap looks even bigger in terms of the momentum-transfer variable t . Nevertheless, the modified SAID solution shown in Figure 5.12 fits the ANKE 1 GeV cross section reduced by 3% and describes also the EDDA data at 1.0144 GeV [62]. Such a 3% reduction in the




FIGURE 5.12: Scaled ANKE proton-proton elastic differential cross sections at 1.0, 2.0, and 2.8 GeV with statistical errors compared to the SAID 2007 solution [9] and a modified (“new”) partial wave solution where the ANKE data have been taken into account. For presentational reasons the 2.0 and 2.8 GeV data and curves have been reduced by factors of 0.5 and 0.25, respectively. The best agreement with the new partial wave data was achieved by scaling the ANKE data with factors 0.97, 0.96, and 1.03 at the three energies. Such factors are within the uncertainties given in Table 5.1.

ANKE normalisation at this energy is consistent with the results of a combined fit of (Equation 5.12) to the EDDA and the Coulomb-corrected ANKE data.

In the forward direction the number of proton-proton elastic scattering amplitudes reduces from five to three and the imaginary parts of these amplitudes are determined completely by the spin-averaged and spin-dependent total cross sections with the help of the generalised optical theorem. The corresponding real parts have been estimated from forward dispersion relations, where these total cross sections provide the necessary input [63]. All the terms contribute positively to the value of $A(GK)$ and, using the optical theorem, the lower bound,

$$A \geq (\sigma_{\text{tot}})^2/16\pi, \quad (5.13)$$

is obtained by taking the pp spin-averaged total cross section σ_{tot} . This lower bound and the full Grein and Kroll estimates for A [63] are both shown in Figure 5.13 where, for consistency, the same values of σ_{tot} were used in the two calculations [57].

T_p GeV	A $\frac{\text{mb}}{(\text{GeV}/c)^2}$	B $(\text{GeV}/c)^{-2}$	C $(\text{GeV}/c)^{-4}$	$A(\text{Corr.})$ $\frac{\text{mb}}{(\text{GeV}/c)^2}$	$A(\text{GK})$ $\frac{\text{mb}}{(\text{GeV}/c)^2}$
1.0	$136.4 \pm 1.3 \pm 3.8$	6.7 ± 0.4	4.0 ± 3.8	136.7 ± 3.8	135.2
1.6	$131.7 \pm 1.9 \pm 4.5$	7.4 ± 0.3	2.7 ± 1.7	131.1 ± 4.5	128.9
1.8	$128.6 \pm 1.7 \pm 4.4$	7.6 ± 0.2	3.4 ± 1.0	127.6 ± 4.3	125.7
2.0	$127.3 \pm 1.7 \pm 4.5$	7.7 ± 0.2	2.5 ± 0.8	124.0 ± 4.3	123.1
2.2	$117.2 \pm 1.8 \pm 3.0$	7.6 ± 0.2	1.4 ± 0.7	113.1 ± 2.9	120.9
2.4	$119.2 \pm 1.8 \pm 3.7$	8.0 ± 0.2	2.7 ± 0.5	112.8 ± 3.5	118.5
2.6	$111.9 \pm 1.7 \pm 3.4$	7.8 ± 0.2	2.0 ± 0.4	108.8 ± 3.3	116.0
2.8	$108.5 \pm 1.8 \pm 2.8$	8.1 ± 0.2	2.4 ± 0.4	105.0 ± 2.7	113.6

TABLE 5.2: Parameters of the fits of Equation 5.12 to the differential cross sections measured in this experiment. In addition to the statistical errors shown, the second uncertainty in the value of A in the second column represents the combined systematic effects summarised in Table 5.1. The corrected values of the forward cross section, $A(\text{Corr.})$, were obtained using the SAID fit discussed in the text, the associated error bars being purely the systematic ones listed in Table 5.1. These values, which were not subjected to the SAID normalisation factors applied in Figure 5.12, may be compared with those of $A(\text{GK})$, which were determined using the Grein and Kroll forward amplitudes [63].

The 992 MeV IKAR data of Figure 5.10 show a significant rise at small $|t|$ that is a reflection of Coulomb distortion of the strong interaction cross section and this was taken into account through the introduction of explicit corrections [58]. The corrected data were then extrapolated to the forward direction ($t = 0$), using a simple exponential function, which would correspond (Equation 5.12) with $C = 0$. The resulting points at all the energies studied are generally about 10% below the Grein and Kroll predictions [63] and would therefore correspond to smaller real parts of the spin-dependent amplitudes. The extrapolation does, of course, depend upon the Coulomb-corrected data following the exponential fit down to $t = 0$.

Though the ANKE data do not probe such small $|t|$ values as IKAR LNPI [58], and are therefore less sensitive to Coulomb distortions, these effects cannot be neglected since they contribute between about 1.5% and 4.5% at 1.0 GeV though less at higher energies.

It is seen in Figure 5.12 that modified SAID solutions describe well the ANKE measurements at three typical energies and the same is true also at the energies not shown. After fitting the ANKE measurements, there is a facility in the SAID




FIGURE 5.13: The predictions of Grein and Kroll [63] for the values of the forward pp elastic differential cross section (solid line), the corresponding lower limit provided by the spin-independent optical theorem (Equation 5.13) being indicated by the dashed line. The extrapolated ANKE data, corresponding to the $A(\text{Corr.})$ parameter of Table 5.2, are shown with their quoted errors by the (blue) circles, whereas the (red) squares are the published IKAR values [58].

program for switching off the Coulomb interaction without adjusting the partial wave amplitudes [9] and this allows a robust extrapolation of the Coulomb-free cross section to the forward direction. The approach has the advantage that it includes some of the minor Coulomb effects that are contained in the SAID program [64][65]. It takes into account the phase variations present in the partial wave analysis and also the deviations from exponential behaviour for very small momentum transfers, $|t| \lesssim m_{\pi^0}^2 = 0.018 \text{ (GeV}/c)^2$, that are linked to pion exchange. The values for $A(\text{Corr.})$ at $t = 0$ produced in this way are given in Table 5.2 and shown in Figure 5.13. The error bars are purely the systematic uncertainties listed in Table 5.1 and any errors in the angular dependence of the SAID predictions are neglected.

The corrections obtained using the SAID program with and without the Coulomb interaction at 1 GeV are a little larger than those found by the LNPI group using an explicit Coulomb formula [58], in part due to the different relative real parts of the pp amplitude in the two calculations.

The agreement of the ANKE data with the theoretical curve in Figure 5.13 is encouraging and would be even slightly better if the normalisation factors found in the fits to the cross sections in Figure 5.12 were implemented. Nevertheless,

the extrapolated values generally fall a little below the predictions at the higher energies.

5.5 Conclusion

In summary, we have measured the differential cross sections for proton-proton elastic scattering at eight energies between 1.0 and 2.8 GeV in a c.m. angular domain between about 12° – 16° to 25° – 30° , depending on the energy. Absolute normalisation of typically 3% were achieved by measuring the energy loss of the beam as it traversed the target. After taking the Coulomb distortions into account, the extrapolations to the forward direction, are broadly compatible with the predictions of forward dispersion relations.

Although our results are completely consistent with ANL measurements at 2.2 and 2.83 GeV [61], the published IKAR values [58] are lower than ours at 1 GeV by about 8%, though this would be reduced to about 5% if one accepts the renormalisation factor from the SAID fit shown in Figure 5.12.




FIGURE 5.14: Energy-dependent phase shift analysis parameters: δ phase shift and ρ ($\cos\rho = \eta$, where η is the absorption parameter) for 1S_0 and 1D_2 partial waves from the 2015 SAID solution [51], in comparison with the older solution from 2007 [8].

The new ANKE data have a significant influence on a partial wave analysis of the pp elastic scattering. In the modified SAID solution, the 1S_0 and 1D_2 waves in particular significantly change at high energies (Figure 5.14). On a more practical level, the measurements will also be a valuable tool in the normalisation of other experiments.

Chapter 6

Analysing power in proton-neutron quasi-free elastic scattering

The analysing power A_y in $\vec{p}n$ quasi-free elastic scattering has been measured at small angles at COSY-ANKE using polarised proton beam at 0.796 GeV and five other beam energies between 1.6 and 2.4 GeV incident on the unpolarised deuteron cluster-jet target. The use of deuterium as an effective neutron target is possible because of the small binding energy of the deuteron (about 2.2 MeV). Consideration of the quasi-free elastic scattering is based on the assumption that the incoming particle is being scattered by only one of two nucleons in the deuteron, while no momentum is transferred to the second nucleon (this nucleon acts as a spectator). In general, the quasi-free scattering is considered to be realized when the momentum transfer from a beam particle to a scattered one (P_t) is large enough as compared with the spectator particle momentum (P_{sp}). Unfortunately, no neutron detection system is available at ANKE-COSY, and hence, no direct measurement of the scattered neutron is possible. The asymmetries of $\vec{p}n$ quasi-free elastic scattering were obtained from the coincidence events, where the fast proton is detected in the ANKE Forward Detector (FD) and the slow spectator proton in a silicon tracking telescope (STT).

This chapter will describe the details of the analysis, unique for the $\vec{p}n$ quasi-free elastic scattering studies at ANKE.

6.1 Beam polarisation

Similar to $\vec{p}p$ elastic scattering experiment, in $\vec{p}d$ measurements the beam polarisation was reversed every subsequent cycle. For each polarisation mode cycles of 180 s or 300 s duration were used, with the last 20 s of each cycle being reserved for the measurement of the beam polarisation with the EDDA detector. The measurement of the proton beam polarisation for the runs with deuteron target followed the same steps as for $\vec{p}p$ ANKE experiment, discussed in Section 4.1.

The weighted averages of the beam polarisations determined over all data at six beam energies are given in Table 6.1. The changes in sign reflect the number of spin flips required to pass through the imperfection resonances, described in detail in Section 3.1.2. The COSY settings for each of the six kinetic energies of the beam were prepared independently and, for this reason, the magnitude of the polarisation may not decrease monotonically with increasing value of the beam energy.

T_p (GeV)	0.796	1.6	1.8	1.965	2.157	2.368
P	0.511	0.378	-0.508	-0.476	-0.513	0.501
ΔP	± 0.001	± 0.001	± 0.003	± 0.005	± 0.005	± 0.004

TABLE 6.1: The values of the mean polarisations P determined with the EDDA polarimeter averaged over all the data at the beam energy T_p . Only statistical errors are given in the table.

The variation of the beam polarisation values among the cycles at any given energy was checked using EDDA with various selections and combinations of the cycles. All the studies yield the consistent results within the uncertainties at any given energy.

6.2 Event selection

Neutron detection is not possible using the ANKE detection system. Therefore, in order to select quasi-free elastic scattering events in the ANKE experiment one proton is measured in the STT in coincidence with the fast proton detected in the FD. The PLUTO simulation [66] has been performed to estimate the rate of pp and pn quasi-free scattering within the acceptance of the ANKE detectors. Figure 6.1 shows the expected counts of $pd \rightarrow ppn_{sp}$ and $pd \rightarrow pnp_{sp}$ reactions at ANKE. The example is shown for beam kinetic energy $T_p = 0.796$ GeV.




FIGURE 6.1: The acceptance $pd \rightarrow ppn_{sp}$ (black histograms) and $pd \rightarrow pnp_{sp}$ (red histograms) reactions simulated in the framework of spectator model at $T_p = 0.796$ GeV. Coincidence events, where one proton is detected in the FD and other one in the STT1 (left panel) or in the STT2 (right panel) are depicted.

The simulation has demonstrated that the pp quasi-elastic scattering is kinematically suppressed in the STT2 due to the FD being an one-arm detector. Therefore, all the coincidence events of the protons detected in the STT2 and the FD are assumed to belong to pn quasi-free elastic scattering. Namely, the proton detected in the STT2 is assumed to be “spectator” one, whereas the particle detected in the FD is regarded as a proton, scattered elastically on a quasi-free neutron.

The proton momentum reconstruction in the STT and the FD followed the same steps as described in Chapter 4. The STTs detect particles with energy threshold of 2 MeV, corresponding to the proton momentum $P_p > 70 \text{ MeV}/c$ [67]. This momentum of spectator proton should be smaller than momentum transfer for the spectator model to be applicable. The careful handling of the data within the spectator model scenario is discussed in greater detail in Section 6.5.

The quasi-free pn elastic scattering events are identified through the evaluation of the missing mass in the $pd \rightarrow ppX$ reaction. As can be seen from the typical example shown in Figure 6.2, at a beam energy of 2.157 GeV, the missing mass peak is well positioned at the neutron mass. The upper limit of the background estimation (details in Section. 6.4) is small compared with the statistical errors of the analysing power.




FIGURE 6.2: Missing-mass $M_X(pp \rightarrow ppX)$ spectra obtained for a beam energy of $T_p = 2.157$ GeV. The peak is consistent with the neutron mass value.

6.3 Asymmetry determination

Since the FD is a single-arm detector, the analysing power for the quasi-free elastic scattering has to be calculated from the simple asymmetries of counts corresponding to different orientations of the beam polarisation. The deuterium cluster target was unfortunately not very stable during this experiment causing large variation of the luminosity (beam-target overlap). Hence, the dedicated normalisation procedure had to be prepared.

In this case, the asymmetry is introduced in terms of normalised numbers of counts for the two orientations of the beam polarisation:

$$\varepsilon(\theta) = \frac{N^\uparrow(\theta)/N_{norm}^\uparrow - N^\downarrow(\theta)/N_{norm}^\downarrow}{N^\uparrow(\theta)/N_{norm}^\uparrow + N^\downarrow(\theta)/N_{norm}^\downarrow}. \quad (6.1)$$

As was discussed in Chapter 5, the normalisation factor must be chosen in such a way, that it does not depend on the beam polarisation. For the given analysis, normalisation factor $N_{norm} = kN_{d1} + N_{d2}$ was constructed from the number of deuterons detected in the STT1 (N_{d1}) and the STT2 (N_{d2}). The polarisation influence can be cancelled in the N_{norm} , if one requires

$$\frac{kN_{d1}^\uparrow - N_{d2}^\uparrow}{kN_{d1}^\uparrow + N_{d2}^\uparrow} = \frac{N_{d2}^\downarrow - kN_{d1}^\downarrow}{N_{d2}^\downarrow + kN_{d1}^\downarrow}. \quad (6.2)$$

The equalising coefficient k is, hence, determined as

$$k = \sqrt{\frac{1}{(N_{d1}^{\uparrow}/N_{d2}^{\uparrow})(N_{d1}^{\downarrow}/N_{d2}^{\downarrow})}} = \sqrt{\frac{N_{d2}^{\downarrow}N_{d2}^{\uparrow}}{N_{d1}^{\uparrow}N_{d1}^{\downarrow}}} = \frac{N_2}{N_1}. \quad (6.3)$$

This approach is based on the several assumptions, including acceptance stability, equality of polarisation values for two modes $|P_{\uparrow}| = |P_{\downarrow}|$, and constant ratio of dead times for the different triggers.

6.3.1 Normalisation via proton-deuteron elastic scattering

It was possible to check the proposed normalisation procedure by comparison with proton-deuteron elastic scattering asymmetry obtained via the cross-ratio method. As was explained in Chapters 2 and 4, the cross-ratio method provides precision results without the first order systematic uncertainties. The cross-ratio asymmetry was measured using stopped deuterons that were identified in the STTs. Similar to $\vec{p}p$ analysis, the angles of the stopped deuterons were deduced from the energy deposits measured in the three layers of the STT rather than from a direct angular measurement.




FIGURE 6.3: Comparison of the $\vec{p}d$ elastic scattering asymmetries $\varepsilon(\theta_{cm})$ at $T_p = 1.965$ GeV obtained in two different ways: blue triangles stand for the cross-ratio method; red squares correspond to the simple left-right asymmetry, using the normalisation procedure described in the text.

We have calculated the left-right asymmetry of the deuterons scattered to the STTs in each polar angular interval as

$$\varepsilon(\theta) = \frac{L(\theta) - R(\theta)}{L(\theta) + R(\theta)} \quad (6.4)$$

from the geometrical means of number of particles scattered in the given polar angle interval to the left L and particles scattered in the same angle interval, but to the right R in respect to the beam polarisation. The results for the asymmetry calculated via the cross-ratio method are compared to the simple left-right asymmetry calculated using the new normalisation procedure in Figure 6.3. The good agreement of the asymmetries shown in Figure 6.3 confirms the validity of the proposed normalisation procedure.

Using the polarisation values from Table 6.1, the analysing power A_y in proton-deuteron elastic scattering was obtained at all six energies. The ANKE data at 0.796 GeV agree with SATURNE measurements [68] within the systematic error bars. Besides providing the normalisation check, these data, summarised in the Figure 6.4 and Appendix D, can be used independently for polarimetry in other experiments.




FIGURE 6.4: Analysing power $A_y(\theta_{cm})$ in $\vec{p}d$ elastic scattering (colored filled circles), along with the existing experimental data from SATURNE (black circles) at $T_p = 0.796$ GeV [68]. Shown values include statistical errors only.

6.4 Background correction

As seen in Figure 6.2, there is some background under neutron peak in the missing mass distribution. In case the background does not depend on the beam polarisation, the true value of the asymmetry ε_{true} can be calculated as

$$\varepsilon_{true} = \varepsilon_{meas}(1 + N_{bg}/N_0) \equiv \varepsilon_{meas}(1 + k_{bg}), \quad (6.5)$$

where N_{bg} is the background count within the chosen range of kinematic variables, and N_0 is the count related to the unpolarised cross section within the same range.




FIGURE 6.5: The difference of normalised missing mass distributions for two polarisation modes at 1.8 GeV.

To check the validity of the background correction via Equation 6.5, the difference of missing mass distributions for different beam polarisation orientations was studied (Figure 6.5). The missing mass spectra for every cycle were normalised by number of elastic deuterons (N_{norm}). Then all normalised distributions obtained at the same beam polarisation mode were summed up with the proper error recalculation. The number of the selected cycles with the beam polarisation up ($\uparrow$) and the beam polarisation down ($\downarrow$) may not necessary be the same. Hence, the further correction for the number of cycles should be undertaken for the final difference of missing mass spectra. As seen in Figure 6.5, contributions of the unpolarised counts are indeed cancelled in such a distribution. The difference of neutron peaks can be fitted by Gaussian, with no background in the vicinity. Similar figures at other energies prove that there is no polarised background and Equation 6.5 can be used for the asymmetry correction.

The true asymmetry should not change among various M_x ranges within the neutron peak, hence it should be possible to roughly estimate the background effect from comparison of the measured asymmetry for different M_x ranges. The large statistical errors, however, do not allow to estimate only very roughly the upper limit of k_{bg} to be 0.05-0.07 within the $\pm 2\sigma$ range.

For more precise estimation on k_{bg} , two background shape hypotheses were investigated: polynomial (Figure 6.6 left panel) and polynomial + Gaussian (Figure 6.6 right panel). The sum of normalised missing mass spectra at 1.8 GeV, fit with Gaussian (centered at neutron mass) and background of different shapes, are shown in Figure 6.6 by black curve. Red dashed curves represent the polynomial approximation for the background, the green dashed curve represents the polynomial + Gaussian (mean value of $m_x \approx 0.86$ GeV/ c^2). The polynomial background does not describe well the tail on the left side of the neutron peak, but the central part of the n-peak is described very well within the $1.5 - 2\sigma$. Hence, even if we consider an additional Gaussian, that helps to describe the left side of the peak, the contribution to the central part is negligible. As a result, both background shapes would give close values of k_{bg} of 0.03 for polynomial, and 0.026 for polynomial + Gaussian shapes.




FIGURE 6.6: The sum of normalised missing mass spectra at 1.8 GeV (blue points), fit with Gaussian and background of different shapes (black curve). Red dashed curves represent the polynomial approximation for the background, the green dashed curve represents the polynomial + Gaussian with $m_x \approx 0.86$ GeV/ c^2 .

6.5 Validity of spectator model

The analysis in the first approximation, when all the coincidence events of protons detected in the FD and in the STT2 are assumed to belong to pn quasi-free elastic

scattering, yields the results that are consistent with the existing SAID solution and experimental data at 0.796 GeV. However, the Figure 6.7 shows that the acceptance of the ANKE detection systems in terms of the P_{sp}/P_t ratio spans from 0.3 up to 1.6, where spectator model is not supposed to be valid. Indeed,




FIGURE 6.7: The ANKE acceptance in terms of the P_{sp}/P_t ratio dependence on the c.m. scattering angle θ_{cm} at the beam energy of $T_p = 0.796$ GeV.

the more careful analysis showed the dependence of the analysing power on the P_{sp}/P_t ratio. It is seen from Figure 6.8 that at least $P_{sp}/P_t < 0.6$ cut is necessary for the analysis at $T_p = 0.796$ GeV. Therefore, the careful handling of the data within the spectator model scenario is necessary.

The following studies on this matter have shown that to separate quasi-free scattering another cut of $P_t < 200$ MeV/c is necessary. The underlying reasons are not entirely clear yet and the investigations are undergoing.

6.5.1 Quasi-free elastic proton-proton scattering

The analysing power in proton-proton elastic scattering offers an interesting possibility for the study of the applicability of the spectator model cuts applied for proton-neutron quasi-free elastic scattering. Namely, we can identify the quasi-free $\vec{p}p$ elastic scattering via detecting the slow proton in the STT and the fast proton in the FD. It should be noted that detecting both scattered protons in the STT and FD allows to get closer to the spectator model, because there is no $P_{sp} < 70$ MeV/c limit anymore. However, applying the same cuts as towards $\vec{p}n$ quasi-free elastic scattering, we can compare the analysing power A_y in quasi-free $\vec{p}p$ to the




FIGURE 6.8: Analysing power A_y dependence on the P_{sp}/P_t ratio at the kinetic beam energy $T_p = 0.796$ GeV. The black horizontal line shows the A_y values expected from SAID 2007 solution.

A_y obtained in free $\vec{p}p$ elastic scattering (Chapter 4). The results are shown in Figure 6.9 at 0.796 GeV (upper panel) and 1.6 GeV (lower panel). At higher energies the A_y values could not be retrieved in the acceptance of the ANKE setup.

The points in red depict the result of the analysis without any cuts on the P_{sp}/P_t ratio. The green triangles are the result of the same cut procedure ($P_{sp}/P_t < 0.5$ and $P_t > 200$ MeV/c) as for pn analysis. Finally, blue empty symbols show the pp elastic data: circles stand for data obtained from STT and squares - from FD.

The analysis of the quasi-free $\vec{p}p$ elastic events proves the justification of the applied cuts within this work. However, the reasons underneath the cuts are not completely understood and are still to be studied in detail. This demonstrates that pn data at 0.796 GeV in the angular range between 18° and 23° indeed correspond to the quasi-free scattering within the applied cuts. The higher energies are influenced much less by the applied cuts, because already for 1.6 GeV the transfer momentum is higher than 180 MeV/c.

6.6 Results and discussion

The results of the A_y measurements at ANKE for pn quasi-elastic scattering are shown in Figure 6.10. The SAID 2007 solution [8], which is only valid up to




FIGURE 6.9: Analysing power in $\bar{p}p$ quasi-free elastic scattering without any cuts (red points), with the same cuts as for $\bar{p}n$ quasi-free elastic scattering (green triangles), compared to the analysing power obtained from free proton-proton elastic scattering at ANKE: FD (blue empty squares) and STT (blue empty circles). The results are shown at $T_p = 0.796$ GeV (upper panel) and $T_p = 1.6$ GeV (lower panel).

1.3 GeV kinetic energy, is shown by the solid black line at the $T_p = 0.796$ GeV in Figure 6.10. This energy was specifically chosen for the comparison with the existing data [69–72] and SAID solution [8]. After the appropriate cuts, that ensure the quasi-free elastic scattering, the analysing power in proton-neutron quasi-free elastic scattering could have been obtained. The agreement between the new ANKE data and other existing data at 0.796 GeV and also with the SAID 2007 prediction is very good.

The higher energies, which were the main purpose of this experiment, have even




FIGURE 6.10: ANKE measurements of the analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-elastic scattering. Upper panel: the ANKE results (red filled circles) at 0.796 GeV are compared with the curves corresponding to the SAID 2007 prediction (solid black line) [9] and other existing measurements (black open symbols) [69–72]. Middle panel: the preliminary SAID solution for the analysing power in pn quasi-free elastic scattering at beam kinetic energy $T_p = 1.6$ GeV is shown in red dashed curve along with the ANKE (red filled circles) and SATURNE (black empty squares) measurements at this energy. Lower panel: ANKE data at four higher energies (filled circles) and the results from Argonne National Laboratory at 2.2 GeV energy [73, 74] (black open symbols). The data at lower panel are scaled for the ease of presentation. Only statistical errors are shown.

better acceptance towards the proton-neutron quasi-free elastic scattering, resulting in the precision data in the non-explored angular and energy region. Results for the higher energies presented in Figure 6.10 were obtained using the data with the cut of $P_{sp}/P_t < 0.4$. This cut does not drastically decrease the available statistics because the FD acceptance for larger P_{sp}/P_t ratio is small. At 1.6 GeV

the loss of statistics due to the limit on the P_{sp}/P_t is about 25%, and it goes down with increasing of beam energy. Excluding the 1.6 GeV beam energy, presented results involve more than 95% of the total available statistics we have collected.

Another confirmation of the high quality of the obtained data is an overlap with the existing data from Argonne National Laboratory (ANL) at 2.2 GeV [73, 74]. The ANKE data points are in a good agreement with ANL data in the overlapping region, and reasonably continue the angular dependence at the smaller angles.

In Figure 6.10 (lower panel) all the results obtained at ANKE at higher energies are summarised. The absolute values are shown at 2.368 GeV, but for clarity of presentation, the other data are scaled up sequentially with the decreasing energy. The scaling factors are given in the legend. The systematic errors, not shown in the figure, include the uncertainties of the determination of the beam polarisation in the EDDA polarimeter.

Using our and other recent data (including WASA-COSY experimental measurements at 1.3 GeV) the SAID group was able to provide the preliminary solution for pn quasi-elastic scattering at 1.6 GeV (shown by red dashed curve in Figure 6.10 middle panel), extending the PWA range from previously published 1.3 GeV.

6.7 Conclusion

We have measured the analysing power in the proton-neutron quasi-free elastic scattering using proton polarised beam at six energies $T_p = 0.796, 1.6, 1.8, 1.965, 2.157, 2.368$ GeV and deuteron unpolarised target. It was possible to show that spectator model could be used in the given experiment.

The ANKE measurements of the analysing power in the proton-neutron quasi-free elastic scattering at beam kinetic energy of 0.796 GeV are consistent with the older experimental data and SAID solution. Furthermore, it was possible to compare the new ANKE measurements with the ANL data at 2.157 GeV, resulting in a good agreement. This further proves the quality of the data at small angles at 5 new energies in the range from 1.6 to 2.4 GeV. This is an important input in the very scarce pn quasi-free elastic scattering data base. Consequently, increasing of the energy range of SAID PWA for pn up to 1.6 GeV has been possible. The shown curve is still only preliminary, the corresponding publication is being prepared for the submission [51]. ANKE data can further aid the SAID analysis by extracting the differential cross-section values in pn quasi-free elastic scattering.

Chapter 7

Summary and outlook

This dissertation presents three sets of nucleon-nucleon (NN) scattering data at small angles, including *precision* data for the analysing power $A_y(\theta)$ and the unpolarised differential cross section $d\sigma/d\Omega(\theta)$ in proton-proton (pp) elastic scattering, and the *first data set* on the analysing power $A_y(\theta)$ in proton-neutron (pn) quasi-free elastic scattering. These studies, motivated by the lack of experimental data for Partial Wave Analysis (PWA) of NN scattering, provide consistent results and form together an important part of the NN studies programme at the Cooler Synchrotron (COSY) in Jülich.

The determination of the analysing power in $\vec{p}p$ elastic scattering, which relied on the use of the EDDA polarimeter for the absolute measurement of the beam polarisation, provided an interesting opportunity for working with the combination of the ANKE Silicon Tracking Telescopes (STT) and Forward Detector (FD) in the study of small angle scattering. The independent analyses of data gathered with the STT and FD, not only yielded great consistency and increased the angular coverage, but also provided a unique opportunity to study possible sources of systematic uncertainties in great detail. It was concluded that the only significant systematic uncertainty is the 3–5% associated with the EDDA polarimeter. The proton beam kinetic energy of 0.796 GeV was specially chosen for the control studies of the analysing power. The ANKE results at this energy are in striking agreement with world data (SATURNE, ANL and others) and also with the SAID 2007 solution. This reinforced our confidence in the results obtained using the ANKE asymmetry measurements together with the EDDA beam polarimetry values.

At five higher energies, $T_p = 1.6, 1.8, 1.965, 2.157$, and 2.368 GeV, the analysing power values were measured for *the first time* at small angles. The comparison with the current SAID analysis showed significant discrepancies. Not only did the ANKE values lie significantly above the SAID prediction, but the angular dependence was also markedly different from that expected. The SAID group subsequently analysed the world data, this time including the ANKE analysing power measurements, and obtained a modified PWA solution that describes the data significantly better. This led to changes in the lower partial waves (e.g., 3F_2 and 3F_4) by up to 60% at the higher energies.

Values of the unpolarised differential cross sections were obtained at eight proton beam energies, namely $T_p = 1.0, 1.6, 1.8, 2.0, 2.2, 2.4, 2.6$, and 2.8 GeV. While the procedure for identifying pp elastic scattering using the FD was very similar to that for the analysing power experiment, the most complex part of the cross section analysis was the determination of the absolute luminosity. An accuracy of about 3% was achieved via the Schottky method, which measured the energy loss of the beam as it traversed the target. After taking Coulomb distortions into account, the values of the cross sections extrapolated to the forward direction are broadly compatible with the predictions of the forward dispersion relations. In addition, our results are completely consistent with ANL measurements at 2.2 and 2.83 GeV.

The precise ANKE values of the observables in pp elastic scattering have provided the SAID data base and PWA with significant new input, resulting in important changes in the low partial waves. Most surprisingly, the 1S_0 and 1D_2 waves changed compared to the SAID 2007 solution.

In order to investigate $\vec{p}n$ quasi-free elastic scattering, experiments with polarised proton beams and an unpolarised deuterium cluster-jet target were carried out at six energies. The polarisation of the proton beam was determined using the EDDA polarimeter while the asymmetry was measured via the coincidence of events where the fast proton was detected in the FD and spectator proton in the STT. It was shown that the spectator model could be used to extract pn quasi-free elastic scattering at all the energies studied.

Quasi-free $\vec{p}p$ elastic scattering could also be identified at $T_p = 0.796$ and 1.6 GeV using the same analysis cuts as for $\vec{p}n$. The agreement of the analysing powers in $\vec{p}p$ quasi-free elastic scattering and free $\vec{p}p$ elastic scattering further confirmed the correctness of the analysis. The procedure for the asymmetry calculation was also checked by comparing the asymmetry obtained for pd elastic scattering with the developed normalisation procedure to that found using the cross-ratio method.

The analysing powers in $\vec{p}d$ elastic scattering can then also be used for polarimetry in other experiments.

The measurements of the analysing power $A_y(\theta)$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 1.6, 1.8, 1.965, 2.157$ and 2.368 GeV were performed for the first time at such small scattering angles. There is also good agreement in the overlap region with the existing experimental data at 0.796 GeV and 2.157 GeV. The SAID group modified the $I = 0$ partial wave analysis, increasing the upper limit of the solution validity from 1.3 GeV up to 1.6 GeV. Nevertheless, more $\vec{p}n$ scattering data are necessary in order to establish a reliable PWA at higher energies. It is planned to extract the differential cross section of pn quasi-free elastic scattering from these ANKE data.

In summary, the results presented in this dissertation are in a good agreement with the existing world data where these exist. However, the ANKE data fill in many holes in the angular and energy coverage of NN elastic scattering observables and these results are having a significant impact on the partial wave analysis.

Appendices

Appendix A

Angular ranges of EDDA semi-rings

Ring Number	$\Delta\theta_{lab}$
14	42.7°... 36.9°
15	39.8°... 34.1°
16	36.9°... 31.5°
17	34.1°... 28.9°
18	31.5°... 26.5°
19	28.9°... 24.2°
20	26.5°... 22.1°
21	24.2°... 20.1°
22	22.1°... 18.3°
23	20.1°... 16.6°
24	18.3°... 15.0°
25	16.6°... 13.6°
26	15.0°... 12.3°
27	13.6°... 11.1°
28	12.3°... 10.1°
29	11.1°... 9.9°

TABLE A.1: Laboratory angle ranges, corresponding to EDDA rings, in the coordinate system, associated with the detector.

Appendix B

Numerical values of A_y in $\vec{p}p$ elastic scattering

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
0.796	6.6	0.1701	0.0112
0.796	7.8	0.1806	0.006
0.796	9.0	0.2108	0.0047
0.796	10.2	0.2326	0.0042
0.796	11.4	0.2505	0.0042
0.796	12.6	0.2682	0.004
0.796	13.8	0.2897	0.004
0.796	15.0	0.3088	0.0038
0.796	16.2	0.3188	0.0038
0.796	17.4	0.3377	0.0038
0.796	18.6	0.3506	0.0038
0.796	19.8	0.3688	0.004
0.796	21.0	0.3789	0.0042
0.796	22.2	0.4022	0.0042
0.796	23.4	0.4049	0.0045
0.796	24.6	0.4152	0.0054
0.796	25.8	0.4336	0.0052
0.796	27.0	0.4414	0.0058
0.796	28.8	0.4466	0.0059
0.796	31.2	0.4651	0.0137

TABLE B.1: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 0.796$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.6	4.2	0.1025	0.0124
1.6	5.4	0.1124	0.0034
1.6	6.6	0.1282	0.0028
1.6	7.8	0.1456	0.0024
1.6	9.0	0.1578	0.0022
1.6	10.2	0.1711	0.002
1.6	11.4	0.1908	0.002
1.6	12.6	0.2044	0.002
1.6	13.8	0.2179	0.002
1.6	15.0	0.2291	0.002
1.6	16.2	0.2457	0.0018
1.6	17.4	0.2528	0.002
1.6	18.6	0.2609	0.002
1.6	19.8	0.2694	0.002
1.6	21.0	0.2836	0.0026
1.6	22.2	0.2878	0.0034
1.6	23.4	0.3016	0.0043
1.6	25.2	0.3081	0.0041

TABLE B.2: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.6$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.8	5.4	0.1106	0.0052
1.8	6.6	0.1278	0.0042
1.8	7.8	0.1405	0.004
1.8	9.0	0.1544	0.0036
1.8	10.2	0.172	0.0034
1.8	11.4	0.182	0.0034
1.8	12.6	0.1964	0.0034
1.8	13.8	0.2122	0.0034
1.8	15.0	0.2166	0.0034
1.8	16.2	0.2297	0.0031
1.8	17.4	0.2426	0.0035
1.8	18.6	0.2473	0.004
1.8	19.8	0.2584	0.0048
1.8	21.0	0.2656	0.0057
1.8	22.2	0.2707	0.0067
1.8	23.4	0.2719	0.0076
1.8	24.6	0.2776	0.0086
1.8	25.8	0.2822	0.0113

TABLE B.3: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.8$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.965	4.2	0.1034	0.0068
1.965	5.4	0.1102	0.0037
1.965	6.6	0.1393	0.0033
1.965	7.8	0.1571	0.0033
1.965	9.0	0.1781	0.003
1.965	10.2	0.1895	0.0028
1.965	11.4	0.1991	0.0028
1.965	12.6	0.2156	0.0028
1.965	13.8	0.2299	0.0026
1.965	15.0	0.2357	0.0026
1.965	16.2	0.2499	0.0026
1.965	17.4	0.2649	0.0026
1.965	18.6	0.2723	0.0028
1.965	19.8	0.2805	0.0035
1.965	21.0	0.2933	0.0042
1.965	22.2	0.2973	0.0047
1.965	23.4	0.3127	0.0056

TABLE B.4: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.965$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.157	4.2	0.085	0.0044
2.157	5.4	0.0984	0.0028
2.157	6.6	0.1235	0.0026
2.157	7.8	0.1325	0.0024
2.157	9.0	0.1484	0.0024
2.157	10.2	0.1632	0.0022
2.157	11.4	0.1726	0.0022
2.157	12.6	0.1868	0.002
2.157	13.8	0.1913	0.002
2.157	15.0	0.2011	0.0022
2.157	16.2	0.2113	0.002
2.157	17.4	0.2201	0.0022
2.157	18.6	0.2304	0.0026
2.157	19.8	0.2342	0.0032
2.157	21.0	0.2446	0.0036
2.157	22.2	0.2508	0.0042
2.157	23.4	0.2544	0.0112

TABLE B.5: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.157$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.368	4.2	0.0757	0.0044
2.368	5.4	0.0911	0.003
2.368	6.6	0.1036	0.0028
2.368	7.8	0.1183	0.0028
2.368	9.0	0.1289	0.0028
2.368	10.2	0.1353	0.0025
2.368	11.4	0.1507	0.0025
2.368	12.6	0.157	0.0023
2.368	13.8	0.1682	0.0025
2.368	15.0	0.1733	0.0023
2.368	16.2	0.1825	0.0023
2.368	17.4	0.1878	0.0028
2.368	18.6	0.1945	0.0035
2.368	19.8	0.2037	0.0039
2.368	21.0	0.2007	0.0044
2.368	22.2	0.2046	0.0099

TABLE B.6: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.368$ GeV, calculated from STT data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
0.796	14.5	0.3065	0.0045
0.796	15.5	0.3241	0.0046
0.796	16.5	0.3331	0.0047
0.796	17.5	0.3445	0.0048
0.796	18.5	0.3659	0.0048
0.796	19.5	0.373	0.0049
0.796	20.5	0.385	0.0051
0.796	21.5	0.3908	0.0052
0.796	22.5	0.4055	0.0056
0.796	23.5	0.4124	0.0078

TABLE B.7: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 0.796$ GeV, calculated from FD data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.6	11.5	0.1873	0.0085
1.6	12.5	0.1988	0.0031
1.6	13.5	0.2101	0.0026
1.6	14.5	0.223	0.0024
1.6	15.5	0.2373	0.0025
1.6	16.5	0.2448	0.0025
1.6	17.5	0.2566	0.0026
1.6	18.5	0.2604	0.0027
1.6	19.5	0.2723	0.0028
1.6	20.5	0.2756	0.003
1.6	21.5	0.2844	0.003
1.6	22.5	0.2941	0.0031
1.6	23.5	0.2991	0.0033
1.6	24.5	0.3012	0.0034
1.6	25.5	0.3076	0.0036
1.6	26.5	0.3168	0.0054

TABLE B.8: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.6$ GeV, calculated from FD data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.8	12.5	0.194	0.006
1.8	13.5	0.2067	0.0048
1.8	14.5	0.2231	0.0045
1.8	15.5	0.2281	0.0045
1.8	16.5	0.2418	0.0045
1.8	17.5	0.2451	0.0046
1.8	18.5	0.2591	0.0047
1.8	19.5	0.2634	0.0048
1.8	20.5	0.269	0.005
1.8	21.5	0.2784	0.0053
1.8	22.5	0.28	0.0057
1.8	23.5	0.2889	0.0063
1.8	24.5	0.2822	0.0068
1.8	25.5	0.2902	0.0069
1.8	26.5	0.2935	0.0072

TABLE B.9: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.8$ GeV, calculated from FD data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.965	12.5	0.2154	0.0062
1.965	13.5	0.2323	0.0034
1.965	14.5	0.2337	0.0028
1.965	15.5	0.2468	0.0028
1.965	16.5	0.256	0.0029
1.965	17.5	0.2636	0.003
1.965	18.5	0.2749	0.0031
1.965	19.5	0.2789	0.0032
1.965	20.5	0.2835	0.0033
1.965	21.5	0.2926	0.0034
1.965	22.5	0.3001	0.0036
1.965	23.5	0.3098	0.004
1.965	24.5	0.3142	0.0041
1.965	25.5	0.3126	0.0042
1.965	26.5	0.3185	0.0045

TABLE B.10: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.965$ GeV, calculated from FD data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.157	13.5	0.1874	0.0026
2.157	14.5	0.1998	0.0021
2.157	15.5	0.2066	0.002
2.157	16.5	0.2137	0.002
2.157	17.5	0.2226	0.0021
2.157	18.5	0.2265	0.0022
2.157	19.5	0.2302	0.0023
2.157	20.5	0.2373	0.0024
2.157	21.5	0.2438	0.0025
2.157	22.5	0.242	0.0026
2.157	23.5	0.2548	0.0028
2.157	24.5	0.2487	0.003
2.157	25.5	0.2559	0.0032
2.157	26.5	0.2637	0.0035
2.157	27.5	0.2639	0.0036

TABLE B.11: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.157$ GeV, calculated from FD data.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.368	13.5	0.1641	0.0033
2.368	14.5	0.1764	0.0024
2.368	15.5	0.1743	0.0023
2.368	16.5	0.1865	0.0023
2.368	17.5	0.1916	0.0024
2.368	18.5	0.1966	0.0025
2.368	19.5	0.2018	0.0026
2.368	20.5	0.206	0.0028
2.368	21.5	0.2066	0.0029
2.368	22.5	0.2157	0.003
2.368	23.5	0.2195	0.0033
2.368	24.5	0.2147	0.0035
2.368	25.5	0.2135	0.0037
2.368	26.5	0.2215	0.004
2.368	27.5	0.2237	0.0043
2.368	28.5	0.2209	0.0046

TABLE B.12: Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.368$ GeV, calculated from FD data.

Appendix C

Numerical values of $d\sigma/d\Omega$ in pp elastic scattering

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
1.0	12.25	17.72	0.02136	118.7
1.0	12.75	17.64	0.02314	118.1
1.0	13.25	17.36	0.02498	116.2
1.0	13.75	16.98	0.02689	113.7
1.0	14.25	16.78	0.02887	112.4
1.0	14.75	16.49	0.03092	110.5
1.0	15.25	16.41	0.03304	109.9
1.0	15.75	16.18	0.03523	108.4
1.0	16.25	15.87	0.03748	106.2
1.0	16.75	15.67	0.03981	105.0
1.0	17.25	15.47	0.04220	103.6
1.0	17.75	15.17	0.04467	101.6

TABLE C.1: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.0$ GeV.

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
1.0	18.25	14.98	0.04720	100.3
1.0	18.75	14.71	0.04979	98.50
1.0	19.25	14.55	0.05246	97.44
1.0	19.75	14.39	0.05519	96.33
1.0	20.25	14.03	0.05799	93.96
1.0	20.75	13.66	0.06086	91.50
1.0	21.25	13.45	0.06380	90.08
1.0	21.75	13.38	0.06680	89.60
1.0	22.25	12.94	0.06986	86.66
1.0	22.75	12.88	0.07300	86.28
1.0	23.25	12.53	0.07620	83.89
1.0	23.75	12.32	0.07946	82.48
1.0	24.25	11.90	0.08279	79.70

TABLE C.2: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.0$ GeV (continuation).

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
1.6	13.75	23.17	0.04302	96.98
1.6	14.25	22.58	0.04619	94.51
1.6	14.75	21.91	0.04947	91.70
1.6	15.25	21.45	0.05286	89.76
1.6	15.75	20.59	0.05636	86.18
1.6	16.25	20.59	0.05997	86.18
1.6	16.75	19.95	0.06370	83.51
1.6	17.25	19.18	0.06753	80.27
1.6	17.75	19.01	0.07147	79.58
1.6	18.25	18.49	0.07551	77.38
1.6	18.75	17.69	0.07967	74.05
1.6	19.25	17.33	0.08394	72.53
1.6	19.75	16.83	0.08831	70.43
1.6	20.25	16.18	0.09279	67.72
1.6	20.75	15.88	0.09738	66.47
1.6	21.25	15.33	0.1021	64.16
1.6	21.75	14.68	0.1069	61.43
1.6	22.25	14.38	0.1118	60.16
1.6	22.75	13.79	0.1168	57.72
1.6	23.25	13.20	0.1219	55.26
1.6	23.75	12.90	0.1271	53.98
1.6	24.25	12.23	0.1325	51.18
1.6	24.75	12.18	0.1379	50.99
1.6	25.25	11.54	0.1434	48.32
1.6	25.75	11.07	0.1491	46.33
1.6	26.25	10.81	0.1548	45.26

TABLE C.3: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.6$ GeV.

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
1.8	14.25	23.43	0.05197	87.16
1.8	14.75	22.97	0.05566	85.47
1.8	15.25	22.17	0.05947	82.47
1.8	15.75	21.62	0.06341	80.42
1.8	16.25	20.78	0.06747	77.31
1.8	16.75	20.39	0.07166	75.86
1.8	17.25	19.83	0.07597	73.78
1.8	17.75	19.24	0.08040	71.56
1.8	18.25	18.56	0.08495	69.05
1.8	18.75	17.96	0.08963	66.81
1.8	19.25	17.40	0.09443	64.73
1.8	19.75	16.70	0.09935	62.14
1.8	20.25	16.04	0.1044	59.67
1.8	20.75	15.83	0.1095	58.90
1.8	21.25	15.07	0.1148	56.08
1.8	21.75	14.36	0.1202	53.43
1.8	22.25	14.21	0.1258	52.87
1.8	22.75	13.59	0.1314	50.56
1.8	23.25	12.84	0.1372	47.77
1.8	23.75	12.34	0.1430	45.89
1.8	24.25	11.95	0.1490	44.45
1.8	24.75	11.60	0.1551	43.16
1.8	25.25	11.05	0.1614	41.09
1.8	25.75	10.48	0.1677	39.00
1.8	26.25	10.18	0.1742	37.86
1.8	26.75	9.678	0.1807	36.01
1.8	27.25	9.457	0.1874	35.18

TABLE C.4: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.8$ GeV.

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
2.0	14.25	24.5061	0.0577396	82.05
2.0	14.75	23.87	0.06184	79.94
2.0	15.25	23.17	0.06608	77.57
2.0	15.75	22.55	0.07045	75.5
2.0	16.25	21.67	0.07497	72.57
2.0	16.75	20.83	0.07962	69.75
2.0	17.25	20.43	0.08441	68.42
2.0	17.75	19.58	0.08933	65.56
2.0	18.25	18.83	0.09439	63.05
2.0	18.75	18.18	0.09959	60.86
2.0	19.25	17.51	0.1049	58.63
2.0	19.75	17.01	0.1104	56.97
2.0	20.25	16.18	0.116.	54.19
2.0	20.75	15.3.	0.1217	51.24
2.0	21.25	15.04	0.1276	50.36
2.0	21.75	14.38	0.1336	48.15
2.0	22.25	13.43	0.1397	44.98
2.0	22.75	13.21	0.146.	44.23
2.0	23.25	12.67	0.1524	42.44
2.0	23.75	12.01	0.1589	40.20
2.0	24.25	11.35	0.1656	38.00
2.0	24.75	10.74	0.1724	35.95
2.0	25.25	10.47	0.1793	35.04
2.0	25.75	9.949	0.1863	33.31
2.0	26.25	9.499	0.1935	31.8.
2.0	26.75	9.036	0.2008	30.26
2.0	27.25	8.484	0.2083	28.41
2.0	27.75	8.287	0.2158	27.75

TABLE C.5: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.0$ GeV

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
2.2	14.75	23.17	0.06802	70.54
2.2	15.25	22.33	0.07269	67.98
2.2	15.75	21.87	0.07750	66.58
2.2	16.25	20.74	0.08247	63.14
2.2	16.75	19.87	0.08758	60.47
2.2	17.25	19.26	0.09285	58.64
2.2	17.75	18.35	0.09826	55.87
2.2	18.25	17.89	0.1038	54.46
2.2	18.75	17.12	0.1095	52.12
2.2	19.25	16.26	0.1154	49.51
2.2	19.75	15.69	0.1214	47.76
2.2	20.25	14.89	0.1276	45.32
2.2	20.75	13.91	0.1339	42.33
2.2	21.25	13.76	0.1403	41.89
2.2	21.75	13.03	0.1470	39.67
2.2	22.25	12.35	0.1537	37.58
2.2	22.75	11.87	0.1606	36.13
2.2	23.25	11.11	0.1676	33.81
2.2	23.75	10.81	0.1748	32.89
2.2	24.25	10.24	0.1821	31.18
2.2	24.75	9.573	0.1896	29.14
2.2	25.25	9.204	0.1972	28.02
2.2	25.75	8.527	0.2050	25.96
2.2	26.25	8.194	0.2129	24.94
2.2	26.75	7.755	0.2209	23.60
2.2	27.25	7.201	0.2291	21.92
2.2	27.75	6.867	0.2374	20.90
2.2	28.25	6.436	0.2459	19.59

TABLE C.6: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.2$ GeV

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
2.4	15.25	23.11	0.0793	64.50
2.4	15.75	21.91	0.0846	61.14
2.4	16.25	21.35	0.08996	59.56
2.4	16.75	20.35	0.09554	56.77
2.4	17.25	19.15	0.1013	53.42
2.4	17.75	18.60	0.1072	51.89
2.4	18.25	17.77	0.1133	49.57
2.4	18.75	17.06	0.1195	47.61
2.4	19.25	16.23	0.1259	45.29
2.4	19.75	15.30	0.1325	42.68
2.4	20.25	14.93	0.1392	41.67
2.4	20.75	14.17	0.1461	39.52
2.4	21.25	13.12	0.1531	36.62
2.4	21.75	12.76	0.1603	35.61
2.4	22.25	11.87	0.1677	33.12
2.4	22.75	11.45	0.1752	31.95
2.4	23.25	10.81	0.1829	30.17
2.4	23.75	10.16	0.1907	28.33
2.4	24.25	9.786	0.1987	27.31
2.4	24.75	9.043	0.2068	25.23
2.4	25.25	8.573	0.2152	23.92
2.4	25.75	8.060	0.2236	22.49
2.4	26.25	7.604	0.2322	21.22
2.4	26.75	7.213	0.2410	20.12
2.4	27.25	6.702	0.2499	18.70
2.4	27.75	6.441	0.2590	17.97
2.4	28.25	5.983	0.2682	16.70
2.4	28.75	5.671	0.2776	15.82
2.4	29.25	5.392	0.2871	15.05

TABLE C.7: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.4$ GeV

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
2.6	15.75	21.58	0.0916	55.57
2.6	16.25	20.63	0.0975	53.12
2.6	16.75	19.82	0.1035	51.06
2.6	17.25	18.88	0.1097	48.64
2.6	17.75	17.61	0.1161	45.36
2.6	18.25	17.24	0.1227	44.40
2.6	18.75	16.35	0.1295	42.11
2.6	19.25	15.56	0.1364	40.08
2.6	19.75	14.83	0.1435	38.20
2.6	20.25	13.85	0.1508	35.66
2.6	20.75	13.41	0.1582	34.54
2.6	21.25	12.50	0.1659	32.18
2.6	21.75	11.79	0.1737	30.36
2.6	22.25	11.26	0.1816	29.00
2.6	22.75	10.45	0.1898	26.92
2.6	23.25	10.04	0.1981	25.86
2.6	23.75	9.354	0.2066	24.09
2.6	24.25	8.864	0.2153	22.83
2.6	24.75	8.381	0.2241	21.59
2.6	25.25	7.735	0.2331	19.92
2.6	25.75	7.377	0.2422	19.00
2.6	26.25	6.740	0.2516	17.36
2.6	26.75	6.518	0.2611	16.79
2.6	27.25	6.064	0.2707	15.62
2.6	27.75	5.608	0.2806	14.44
2.6	28.25	5.324	0.2906	13.71
2.6	28.75	4.88	0.3007	12.57
2.6	29.25	4.615	0.311	11.89
2.6	29.75	4.348	0.3215	11.20

TABLE C.8: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.6$ GeV

T_p [GeV]	θ_{cm} [°]	$d\sigma/d\Omega(\theta_{cm})$ [mb]	$-t$ (GeV/c) ²	$d\sigma/dt$ [mb/(GeV/c) ²]
2.8	16.25	20.12	0.1049	48.13
2.8	16.75	19.10	0.1115	45.67
2.8	17.25	17.84	0.1182	42.66
2.8	17.75	17.11	0.1251	40.93
2.8	18.25	16.16	0.1321	38.65
2.8	18.75	15.48	0.1394	37.01
2.8	19.25	14.56	0.1469	34.83
2.8	19.75	13.71	0.1545	32.80
2.8	20.25	13.07	0.1624	31.26
2.8	20.75	12.17	0.1704	29.11
2.8	21.25	11.81	0.1786	28.24
2.8	21.75	10.89	0.1870	26.04
2.8	22.25	10.09	0.1956	24.13
2.8	22.75	9.771	0.2044	23.37
2.8	23.25	8.991	0.2133	21.50
2.8	23.75	8.505	0.2225	20.34
2.8	24.25	7.938	0.2318	18.98
2.8	24.75	7.452	0.2413	17.82
2.8	25.25	7.020	0.2510	16.79
2.8	25.75	6.449	0.2609	15.42
2.8	26.25	6.015	0.2709	14.39
2.8	26.75	5.586	0.2812	13.36
2.8	27.25	5.278	0.2916	12.62
2.8	27.75	4.915	0.3022	11.76
2.8	28.25	4.633	0.3129	11.08
2.8	28.75	4.251	0.3239	10.17
2.8	29.25	3.900	0.3350	9.326
2.8	29.75	3.749	0.3463	8.966
2.8	30.25	3.425	0.3577	8.191

TABLE C.9: Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.8$ GeV

Appendix D

Numerical values of A_y in $\vec{p}d$ elastic scattering

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
0.796	7.5	0.2573	0.0116
0.796	8.5	0.2680	0.0040
0.796	9.5	0.2929	0.0032
0.796	10.5	0.3156	0.0029
0.796	11.5	0.3401	0.0029
0.796	12.5	0.3582	0.0028
0.796	13.5	0.3778	0.0024
0.796	14.5	0.3923	0.0027
0.796	15.5	0.4099	0.0029
0.796	16.5	0.4153	0.0031
0.796	17.5	0.4226	0.0034
0.796	18.5	0.4253	0.0037
0.796	19.5	0.4272	0.0040
0.796	20.5	0.4335	0.0044
0.796	21.5	0.4340	0.0049
0.796	22.5	0.4304	0.0057
0.796	23.5	0.4252	0.0066
0.796	24.5	0.4112	0.0076
0.796	25.5	0.4054	0.0090
0.796	26.5	0.4242	0.0110
0.796	27.5	0.3984	0.0164
0.796	28.5	0.4055	0.0904

TABLE D.1: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 0.796$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.6	5.5	0.1153	0.0031
1.6	6.5	0.1056	0.0016
1.6	7.5	0.1267	0.0013
1.6	8.5	0.1390	0.0014
1.6	9.5	0.1463	0.0013
1.6	10.5	0.1561	0.0015
1.6	11.5	0.1674	0.0016
1.6	12.5	0.1694	0.0018
1.6	13.5	0.1790	0.0021
1.6	14.5	0.1815	0.0024
1.6	15.5	0.1736	0.0027
1.6	16.5	0.1840	0.0032
1.6	17.5	0.1719	0.0038
1.6	18.5	0.1877	0.0046
1.6	19.5	0.1925	0.0100

TABLE D.2: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.6$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.8	5.5	0.1227	0.0017
1.8	6.5	0.1324	0.0012
1.8	7.5	0.1430	0.0010
1.8	8.5	0.1538	0.0009
1.8	9.5	0.1647	0.0009
1.8	10.5	0.1807	0.0011
1.8	11.5	0.1855	0.0012
1.8	12.5	0.1909	0.0014
1.8	13.5	0.1997	0.0016
1.8	14.5	0.2067	0.0019
1.8	15.5	0.2018	0.0022
1.8	16.5	0.1969	0.0027
1.8	17.5	0.1970	0.0037

TABLE D.3: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.8$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.965	4.5	0.1003	0.0110
1.965	5.5	0.0968	0.0024
1.965	6.5	0.1077	0.0019
1.965	7.5	0.1230	0.0017
1.965	8.5	0.1324	0.0014
1.965	9.5	0.1410	0.0016
1.965	10.5	0.1524	0.0018
1.965	11.5	0.1616	0.0020
1.965	12.5	0.1645	0.0023
1.965	13.5	0.1665	0.0027
1.965	14.5	0.1738	0.0032
1.965	15.5	0.1601	0.0039
1.965	16.5	0.1620	0.0049
1.965	17.5	0.1560	0.0127

TABLE D.4: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.965$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.157	4.5	0.0862	0.0074
2.157	5.5	0.0922	0.0028
2.157	6.5	0.1053	0.0024
2.157	7.5	0.1127	0.0020
2.157	8.5	0.1264	0.0019
2.157	9.5	0.1330	0.0022
2.157	10.5	0.1415	0.0025
2.157	11.5	0.1487	0.0029
2.157	12.5	0.1550	0.0033
2.157	13.5	0.1606	0.0039
2.157	14.5	0.1603	0.0047
2.157	15.5	0.1532	0.0059
2.157	16.5	0.1500	0.0107

TABLE D.5: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 2.157$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.368	4.5	0.0821	0.0045
2.368	5.5	0.0910	0.0023
2.368	6.5	0.0993	0.0021
2.368	7.5	0.1132	0.0017
2.368	8.5	0.1179	0.0018
2.368	9.5	0.1248	0.0020
2.368	10.5	0.1319	0.0024
2.368	11.5	0.1355	0.0028
2.368	12.5	0.1430	0.0033
2.368	13.5	0.1474	0.0040
2.368	14.5	0.1357	0.0050
2.368	15.5	0.1333	0.0082
2.368	16.5	0.1498	0.0107

TABLE D.6: Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 2.368$ GeV.

Appendix E

Numerical values of A_y in $\vec{p}n$ quasi-free elastic scattering

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
0.796	18	0.270	0.018
0.796	20	0.290	0.018
0.796	22	0.278	0.016
0.796	24	0.328	0.022

TABLE E.1: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 0.796$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.6	14	0.137	0.018
1.6	17	0.151	0.015
1.6	20	0.153	0.015
1.6	23	0.170	0.016
1.6	26	0.169	0.018

TABLE E.2: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 1.6$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.8	13.5	0.125	0.011
1.8	16.5	0.145	0.008
1.8	19.5	0.152	0.008
1.8	22.5	0.148	0.009
1.8	25.5	0.158	0.010

TABLE E.3: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 1.8$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
1.965	13.5	0.108	0.016
1.965	16.5	0.119	0.011
1.965	19.5	0.138	0.011
1.965	22.5	0.112	0.012
1.965	25.5	0.141	0.013
1.965	28.5	0.127	0.022

TABLE E.4: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 1.965$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.157	14	0.091	0.016
2.157	17	0.103	0.013
2.157	20	0.111	0.013
2.157	23	0.143	0.014
2.157	26	0.115	0.017
2.157	29	0.141	0.026

TABLE E.5: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 2157$ GeV.

T_p [GeV]	θ_{cm} [°]	$A_y(\theta_{cm})$	$\Delta A_y(\theta_{cm})$
2.368	17	0.085	0.011
2.368	20	0.102	0.012
2.368	23	0.102	0.013
2.368	26	0.122	0.016
2.368	29	0.100	0.021

TABLE E.6: Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p = 2.368$ GeV.

List of Figures

1.1	The Standard Model of elementary particles (schematic depiction), with the three generations of matter in the first three columns, gauge bosons in the fourth column, and the Higgs boson in the fifth [1].	2
1.2	The value of the "running" coupling constant α_s , as a function of the energy scale E . The curve that slopes downwards is a prediction of the asymptotic freedom in QCD, while the empty circles show the measurements that have been made [7].	4
1.3	Abundance plots of c.m. scattering angle (θ_{cm}) versus beam kinetic energy (T_{lab}) for experiments on the analysing power A_y (left) and for cross-section $d\sigma/d\Omega$ (right) in proton-proton elastic scattering [15].	6
1.4	Abundance plot of c.m. scattering angle (θ_{cm}) versus beam kinetic energy (T_{lab}) for experiments on the analysing power A_y in proton-neutron scattering [15].	7
2.1	Madison convention for the definition of the coordinate system in the polarised experiments. The z axis is along the incident beam momentum, while the scattering is in the xz plane. The spin quantization axis $\mathbf{S}$ is defined in terms of β , the angle between $\mathbf{S}$ and beam direction, z -axis, and ϕ , the angle between its projection on xy plane and the y axis.	12
2.2	The idealistic symmetric arrangement of the two-detector system.. The two cases of beam polarisation are shown: "up" in red and "down" in green. The corresponding counts of particles scattered to the detectors are described in text.	14
3.1	The COSY accelerator facility at Forschungszentrum Jülich. The positions of the ANKE spectrometer and the EDDA polarimeter at the ring are shown. In the bottom of the figure the sources providing polarised and unpolarised protons (or deuterons), the cyclotron that accelerates the particles to the injection energy and the Low Energy (LE) Polarimeter can be seen.	18
3.2	Setup of the polarised ion source at COSY [28]. The negatively charged ions are produced in a charge exchange process between a neutral nuclear polarised hydrogen beam and a fast neutral cesium beam.	19
3.3	The schematic layout of the full EDDA detector setup (left) and photo (right) of the EDDA polarimeter.	23

3.4	The ANKE spectrometer setup (top view), showing the positions of the hydrogen cluster-jet target, the silicon tracking telescopes (STT), and the forward detector (FD).	24
3.5	The cluster-jet target installed at ANKE.	25
3.6	Cluster production process in a Laval-nozzle. The cluster-jet beam is extracted and shaped using the skimmer. Note the exaggerated scale, the opening of the skimmer is only 700 μm	26
3.7	Photo of a Silicon Tracking Telescope (STT), including the cooling system and read-out electronics.	28
4.1	The polarisation values, calculated ring by ring, are shown versus corresponding laboratory polar angles (according to Appendix A). The sample plots are shown for beam kinetic energies $T_p = 1.8$ GeV and $T_p = 2.157$ GeV.	33
4.2	Difference between the directly measured angle of the track θ and the angle reconstructed from the energy $\theta(E)$. The example plot is shown at $T_p = 2.368$ GeV.	35
4.3	Missing mass $M_X(pp \rightarrow pX)$ spectrum obtained for the beam energy of $T_p = 1.6$ GeV, showing the clear proton peak when detecting one proton in the STT.	36
4.4	Missing mass $M_X(pp \rightarrow pX)$ spectrum obtained from the particles detected in the forward detector at the beam kinetic energy of 1.6 GeV.	38
4.5	Forward detector angular acceptance for $pp \rightarrow pp$ at the beam kinetic energy of 1.6 GeV.	38
4.6	Relative luminosities obtained with various angular cuts (shown at horizontal axis) at $T_p = 1.8$ GeV. Only a small section of the vertical scale is shown to emphasise the small differences.	39
4.7	Difference between the scattering angles reconstructed using the FD θ_{cm}^{FD} and STT θ_{cm}^{STT} . Example plots are shown for $T_p = 0.796$ GeV (left panel) and $T_p = 2.157$ GeV (right panel).	41
4.8	Systematic error due to the possible maximum shift of the scattering angle. Example plots are shown for the beam kinetic energies $T_p = 0.796$ GeV (left panel) and $T_p = 2.368$ GeV (right panel).	42
4.9	The angular dependences of the instability factor $r_{\uparrow}/r_{\downarrow}$ at the beam energies of $T_p = 0.796$ GeV (left panel) and $T_p = 1.8$ GeV (right panel).	43
4.10	Efficiencies of the elastic event reconstruction using the FD in the cycles with the beam polarisation mode up (left panel) and beam polarisation mode down (right panel).	44

4.11	Comparison of the ANKE measurements of the proton analysing power in $\vec{p}p$ elastic scattering using the STT (red filled circles) and the FD (blue filled triangles) systems with the curves corresponding to the SAID 2007 (solid black line) [8] and the revised fit (dashed red) solutions. Only statistical errors are shown so that the systematic uncertainties arising, for example, from the calibration of the EDDA polarimeter have not been included. Also shown are selected results from EDDA (black crosses) [13] at the energies different by no more than 7 MeV and, at 0.796 GeV, LAMPF [46–48], and SATURNE [49] (black open symbols).	45
4.12	Energy-dependent phase shift analysis parameters: δ phase shift and ρ ($\cos\rho = \eta$, where η is the absorption parameter) for 3F_2 and 3F_4 partial waves from the 2014 SAID solution, in comparison with the older solution from 2007.	46
5.1	Beam Current Transformer (BCT) typical raw signal, recorded during the experiment.	51
5.2	BCT signal vs. current I in the calibration wire.	51
5.3	Schottky power spectra for $T_p = 1.0$ GeV (a) and $T_p = 2.0$ GeV (b) obtained during one 300 s cycle and scaled to harmonic number 1. The mean frequencies are indicated by the vertical (red) lines. . . .	54
5.4	Typical mean frequency shift derived from the Schottky power spectra of the type illustrated in Figure 5.3 for $T_p = 1.0$ GeV (left panel) and $T_p = 2.0$ GeV (right panel).	55
5.5	Average frequency shifts within the cycle duration, measured by the new (blue) and old (green) spectrum analysers at $T_p = 1.0$ and $T_p = 2.0$ GeV.	55
5.6	Example of frequency shift caused by the background when the target is on $(\frac{df}{dt})_{ANKE+COSY}$ (left) and when the target is off $(\frac{df}{dt})_{COSY}$ (right) compared to the total frequency shift $(\frac{df}{dt})_{tot}$ with the beam incident on the target (middle).	56
5.7	Variation of the relative change in the mean beam revolution frequency $\Delta f/f_0$ with the relative change in the field strength in the bending magnets $\Delta B/B_0$. The fit defines the momentum compaction factor α . The example is shown for $T_p = 2.0$ GeV.	57
5.8	The luminosity measured in every cycle during the experiment. . . .	58
5.9	Combined ANKE data set of differential cross sections with respect to the four-momentum transfer t compared to fits made on the basis of Equation 5.12. Systematic errors are not shown. The correct values are shown at 1.0 GeV but, for clarity of presentation, the other data are scaled down sequentially in energy by factors of 1.2. The true numerical values of the cross section and fit parameters are given in Appendix C and Table 5.2, accordingly.	61

5.10	Differential cross section for pp elastic scattering. The ANKE data at $T_p = 1$ GeV with statistical errors (blue squares) are compared to the IKAR (LNPI) hydrogen data at 992 MeV (green circles) [58] scaled by a factor of 1.085 and methane results at 991 MeV (red triangles) [60] scaled by a factor of 1.04. At very small values of $ t $ there is a rise caused mainly by Coulomb-nuclear interference.	62
5.11	The ANKE pp differential cross section data at 2.2 GeV (closed blue circles) and 2.8 GeV (closed blue triangles) compared to the ANL results [61] at 2.2 GeV (open red circles) and 2.83 GeV (open red triangles). Systematic errors are not shown. For presentational purposes, both higher energy data sets have been scaled downwards by a common factor of 1.5.	63
5.12	Scaled ANKE proton-proton elastic differential cross sections at 1.0, 2.0, and 2.8 GeV with statistical errors compared to the SAID 2007 solution [9] and a modified (“new”) partial wave solution where the ANKE data have been taken into account. For presentational reasons the 2.0 and 2.8 GeV data and curves have been reduced by factors of 0.5 and 0.25, respectively. The best agreement with the new partial wave data was achieved by scaling the ANKE data with factors 0.97, 0.96, and 1.03 at the three energies. Such factors are within the uncertainties given in Table 5.1.	64
5.13	The predictions of Grein and Kroll [63] for the values of the forward pp elastic differential cross section (solid line), the corresponding lower limit provided by the spin-independent optical theorem (Equation 5.13) being indicated by the dashed line. The extrapolated ANKE data, corresponding to the $A(\text{Corr.})$ parameter of Table 5.2, are shown with their quoted errors by the (blue) circles, whereas the (red) squares are the published IKAR values [58].	66
5.14	Energy-dependent phase shift analysis parameters: δ phase shift and ρ ($\cos\rho = \eta$, where η is the absorption parameter) for 1S_0 and 1D_2 partial waves from the 2015 SAID solution [51], in comparison with the older solution from 2007 [8].	67
6.1	The acceptance $pd \rightarrow ppn_{sp}$ (black histograms) and $pd \rightarrow pnp_{sp}$ (red histograms) reactions simulated in the framework of spectator model at $T_p = 0.796$ GeV. Coincidence events, where one proton is detected in the FD and other one in the STT1 (left panel) or in the STT2 (right panel) are depicted.	71
6.2	Missing-mass $M_X(pp \rightarrow ppX)$ spectra obtained for a beam energy of $T_p = 2.157$ GeV. The peak is consistent with the neutron mass value.	72
6.3	Comparison of the $\vec{p}d$ elastic scattering asymmetries $\varepsilon(\theta_{cm})$ at $T_p = 1.965$ GeV obtained in two different ways: blue triangles stand for the cross-ratio method; red squares correspond to the simple left-right asymmetry, using the normalisation procedure described in the text.	73

6.4	Analysing power $A_y(\theta_{cm})$ in $\bar{p}d$ elastic scattering (colored filled circles), along with the existing experimental data from SATURNE (black circles) at $T_p = 0.796$ GeV [68]. Shown values include statistical errors only.	74
6.5	The difference of normalised missing mass distributions for two polarisation modes at 1.8 GeV.	75
6.6	The sum of normalised missing mass spectra at 1.8 GeV (blue points), fit with Gaussian and background of different shapes (black curve). Red dashed curves represent the polynomial approximation for the background, the green dashed curve represents the polynomial + Gaussian with $m_x \approx 0.86$ GeV/ c^2	76
6.7	The ANKE acceptance in terms of the P_{sp}/P_t ratio dependence on the c.m. scattering angle θ_{cm} at the beam energy of $T_p = 0.796$ GeV.	77
6.8	Analysing power A_y dependence on the P_{sp}/P_t ratio at the kinetic beam energy $T_p = 0.796$ GeV. The black horizontal line shows the A_y values expected from SAID 2007 solution.	78
6.9	Analysing power in $\bar{p}p$ quasi-free elastic scattering without any cuts (red points), with the same cuts as for $\bar{p}n$ quasi-free elastic scattering (green triangles), compared to the analysing power obtained from free proton-proton elastic scattering at ANKE: FD (blue empty squares) and STT (blue empty circles). The results are shown at $T_p = 0.796$ GeV (upper panel) and $T_p = 1.6$ GeV (lower panel).	79
6.10	ANKE measurements of the analysing power $A_y(\theta_{cm})$ in $\bar{p}n$ quasi-elastic scattering. Upper panel: the ANKE results (red filled circles) at 0.796 GeV are compared with the curves corresponding to the SAID 2007 prediction (solid black line) [9] and other existing measurements (black open symbols) [69–72]. Middle panel: the preliminary SAID solution for the analysing power in pn quasi-free elastic scattering at beam kinetic energy $T_p = 1.6$ GeV is shown in red dashed curve along with the ANKE (red filled circles) and SATURNE (black empty squares) measurements at this energy. Lower panel: ANKE data at four higher energies (filled circles) and the results from Argonne National Laboratory at 2.2 GeV energy [73, 74] (black open symbols). The data at lower panel are scaled for the ease of presentation. Only statistical errors are shown.	80

List of Tables

3.1	Resonance strength ϵ_r and the ratio of preserved polarisation P_f/P_i at imperfection resonances for a typical vertical orbit deviation y_{co}^{rms} , without considering synchrotron oscillation.	21
3.2	Resonance strength ϵ_r of intrinsic resonances for a normalized emittance of 1π mm mrad and vertical betatron tune of $Q_y = 3.61$ for different superperiodicities P	22
4.1	The values of the mean polarisations P determined with the EDDA polarimeter averaged over all the data at the beam energy T_p . Only statistical errors are given in the table.	33
4.2	The normalisation factors N obtained in a partial wave fit [51] to the current STT data.	46
5.1	Percentage contributions to the total systematic uncertainty at different proton beam energies T_p . E_1 reflects the statistical and systematic effects in the determination of the Schottky η parameter. E_2 arises from the rest gas effect (including direct measurement errors as well as possible instabilities). E_3 is a measure of the density instability through the 300 s cycle. These contributions have been added in quadrature together with the accuracy of the stopping powers and the precision of the FD analysis to give the total percentage uncertainty in the last column.	60
5.2	Parameters of the fits of Equation 5.12 to the differential cross sections measured in this experiment. In addition to the statistical errors shown, the second uncertainty in the value of A in the second column represents the combined systematic effects summarised in Table 5.1. The corrected values of the forward cross section, $A(\text{Corr.})$, were obtained using the SAID fit discussed in the text, the associated error bars being purely the systematic ones listed in Table 5.1. These values, which were not subjected to the SAID normalisation factors applied in Figure 5.12, may be compared with those of $A(\text{GK})$, which were determined using the Grein and Kroll forward amplitudes [63].	65
6.1	The values of the mean polarisations P determined with the EDDA polarimeter averaged over all the data at the beam energy T_p . Only statistical errors are given in the table.	70

A.1	Laboratory angle ranges, corresponding to EDDA rings, in the co-ordinate system, associated with the detector.	89
B.1	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 0.796$ GeV, calculated from STT data.	91
B.2	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.6$ GeV, calculated from STT data.	92
B.3	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.8$ GeV, calculated from STT data.	92
B.4	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.965$ GeV, calculated from STT data.	93
B.5	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.157$ GeV, calculated from STT data.	93
B.6	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.368$ GeV, calculated from STT data.	94
B.7	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 0.796$ GeV, calculated from FD data.	94
B.8	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.6$ GeV, calculated from FD data.	95
B.9	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.8$ GeV, calculated from FD data.	95
B.10	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 1.965$ GeV, calculated from FD data.	96
B.11	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.157$ GeV, calculated from FD data.	96
B.12	Analysing power A_y in $\vec{p}p$ elastic scattering at $T_p = 2.368$ GeV, calculated from FD data.	97
C.1	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.0$ GeV.	99
C.2	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.0$ GeV (continuation).	100
C.3	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.6$ GeV.	101
C.4	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 1.8$ GeV.	102
C.5	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.0$ GeV	103
C.6	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.2$ GeV	104
C.7	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.4$ GeV	105
C.8	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.6$ GeV	106
C.9	Unpolarised differential cross section $d\sigma/d\Omega$ in pp elastic scattering at $T_p = 2.8$ GeV	107

D.1	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 0.796$ GeV. . . .	109
D.2	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.6$ GeV. . . .	110
D.3	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.8$ GeV. . . .	110
D.4	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 1.965$ GeV. . . .	111
D.5	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 2.157$ GeV. . . .	111
D.6	Analysing power A_y in $\vec{p}d$ elastic scattering at $T_p = 2.368$ GeV. . . .	112
E.1	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 0.796 GeV.	113
E.2	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 1.6 GeV.	113
E.3	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 1.8 GeV.	114
E.4	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 1.965 GeV.	114
E.5	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 2157 GeV.	114
E.6	Analysing power $A_y(\theta_{cm})$ in $\vec{p}n$ quasi-free elastic scattering at $T_p =$ 2.368GeV.	114

Bibliography

- [1] The Standard Model of Elementary Particles. URL https://en.wikipedia.org/wiki/Standard_Model#/media/File:Standard_Model_of_Elementary_Particles.svg.
- [2] M. Gell-Mann. A schematic model of baryons and mesons. *Physics Letters*, 8(3):214 – 215, 1964. ISSN 0031-9163. doi: [http://dx.doi.org/10.1016/S0031-9163\(64\)92001-3](http://dx.doi.org/10.1016/S0031-9163(64)92001-3). URL <http://www.sciencedirect.com/science/article/pii/S0031916364920013>.
- [3] G. Zweig. An SU(3) model for strong interaction symmetry and its breaking. Version 1. 1964.
- [4] E. D. Bloom, D. H. Coward, H. DeStaebler, J. Drees, G. Miller, L. W. Mo, R. E. Taylor, M. Breidenbach, J. I. Friedman, G. C. Hartmann, and H. W. Kendall. High-energy inelastic $e - p$ scattering at 6° and 10° . *Phys. Rev. Lett.*, 23:930–934, Oct 1969. doi: 10.1103/PhysRevLett.23.930. URL <http://link.aps.org/doi/10.1103/PhysRevLett.23.930>.
- [5] M. Breidenbach, J. I. Friedman, H. W. Kendall, E. D. Bloom, D. H. Coward, H. DeStaebler, J. Drees, L. W. Mo, and R. E. Taylor. Observed behavior of highly inelastic electron-proton scattering. *Phys. Rev. Lett.*, 23:935–939, Oct 1969. doi: 10.1103/PhysRevLett.23.935. URL <http://link.aps.org/doi/10.1103/PhysRevLett.23.935>.
- [6] Siegfried Bethke. The 2009 world average of α_s . *The European Physical Journal C*, 64(4):689, 2009. URL <http://link.springer.com/article/10.1140%2Fepjc%2Fs10052-009-1173-1>.
- [7] About the nobel prize in physics 2004. popular information. URL http://www.nobelprize.org/nobel_prizes/physics/laureates/2004/phy2highen.jpg.

- [8] R. A. Arndt, W. J. Briscoe, I. I. Strakovsky, and R. L. Workman. Updated analysis of nn elastic scattering to 3 GeV. *Phys. Rev. C*, 76:025209, Aug 2007. doi: 10.1103/PhysRevC.76.025209. URL <http://link.aps.org/doi/10.1103/PhysRevC.76.025209>.
- [9] Richard A. Arndt, Igor I. Strakovsky, and Ron L. Workman. Nucleon-nucleon elastic scattering to 3 GeV. *Phys. Rev. C*, 62:034005, Aug 2000. doi: 10.1103/PhysRevC.62.034005. URL <http://link.aps.org/doi/10.1103/PhysRevC.62.034005>.
- [10] Scattering Analysis Interactive Dial-in. URL <http://gwdac.phys.gwu.edu>.
- [11] D. Albers, J. Bisplinghoff, R. Bollmann, K. Büßer, P. Cloth, R. Daniel, O. Diehl, F. Dohrmann, H. P. Engelhardt, J. Ernst, P. D. Eversheim, M. Gasthuber, R. Gebel, J. Greiff, A. Groß, R. Groß-Hardt, S. Heider, A. Heine, F. Hinterberger, M. Igelbrink, R. Jahn, M. Jeske, U. Lahr, R. Langkau, J. Lindlein, R. Maier, R. Maschuw, T. Mayer-Kuckuk, F. Mosel, M. Müller, M. Münstermann, D. Prasuhn, H. Rohdjeß, D. Rosendaal, U. Roß, P. von Rossen, H. Scheid, N. Schirm, M. Schulz-Rojahn, F. Schwandt, V. Schwarz, W. Scobel, G. Sterzenbach, H. J. Trelle, A. Wellinghausen, W. Wiedmann, K. Woller, and R. Ziegler. Proton-proton elastic scattering excitation functions at intermediate energies. *Phys. Rev. Lett.*, 78:1652–1655, Mar 1997. doi: 10.1103/PhysRevLett.78.1652. URL <http://link.aps.org/doi/10.1103/PhysRevLett.78.1652>.
- [12] M. Altmeier, F. Bauer, Bisplinghoff, et al. Excitation functions of the analyzing power in pp scattering from 0.45 to 2.5 GeV. *Phys. Rev. Lett.*, 85:1819–1822, Aug 2000. doi: 10.1103/PhysRevLett.85.1819. URL <http://link.aps.org/doi/10.1103/PhysRevLett.85.1819>.
- [13] M. Altmeier, F. Bauer, J. Bisplinghoff, K. Büßer, M. Busch, T. Colberg, L. Demirörs, H.P. Engelhardt, P.D. Eversheim, K.O. Eyser, O. Felden, R. Gebel, M. Glende, J. Greiff, F. Hinterberger, E. Jonas, H. Krause, T. Lindemann, J. Lindlein, B. Lorentz, R. Maier, R. Maschuw, A. Meinerzhagen, D. Prasuhn, H. Rohdjeß, D. Rosendaal, P. von Rossen, N. Schirm, V. Schwarz, W. Scobel, H.-J. Trelle, K. Ulbrich, E. Weise, A. Wellinghausen, and R. Ziegler. Excitation functions of the analyzing power in elastic proton-proton scattering from 0.45 to 2.5 GeV. *The European Physical Journal A - Hadrons and Nuclei*, 23(2):351–364, 2005. ISSN 1434-6001. doi: 10.1140/epja/i2004-10081-1. URL <http://dx.doi.org/10.1140/epja/i2004-10081-1>.

- [14] F. Bauer, J. Bisplinghoff, K. Büßer, M. Busch, T. Colberg, L. Demirörs, C. Dahl, P. D. Eversheim, O. Eyser, O. Felden, R. Gebel, J. Greiff, F. Hinterberger, E. Jonas, H. Krause, C. Lehmann, J. Lindlein, R. Maier, A. Meinerzhagen, C. Pauly, D. Prasuhn, H. Rohdjeß, D. Rosendaal, P. von Rossen, N. Schirm, W. Scobel, K. Ulbrich, E. Weise, T. Wolf, and R. Ziegler. Measurement of spin-correlation parameters A_{NN} , A_{SS} , and A_{SL} at 2.1 GeV in proton-proton elastic scattering. *Phys. Rev. Lett.*, 90:142301, Apr 2003. doi: 10.1103/PhysRevLett.90.142301. URL <http://link.aps.org/doi/10.1103/PhysRevLett.90.142301>.
- [15] NN-OnLine: Abundance plots of the NN database. URL <http://nn-online.org/NN>.
- [16] J. Bystricky, F. Lehar, and P. Winternitz. Formalism of nucleon-nucleon elastic scattering experiments. *Journal de Physique*, 39(1):1–32, 1978. doi: 10.1051/jphys:019780039010100. URL <https://hal.archives-ouvertes.fr/jpa-00208735>.
- [17] Catherine Lechanoine-LeLuc and Francois Lehar. Nucleon-nucleon elastic scattering and total cross sections. *Rev. Mod. Phys.*, 65:47–86, Jan 1993. doi: 10.1103/RevModPhys.65.47. URL <http://link.aps.org/doi/10.1103/RevModPhys.65.47>.
- [18] Hans Paetz gen. Schieck. *Nuclear Reactions. An Introduction*, volume 882. Springer-Verlag Berlin Heidelberg, 2014. doi: 10.1007/978-3-642-53986-2. URL <http://www.springer.com/gp/book/9783642539855>.
- [19] P. Huber and K.P. Meyer. *Proceedings of the 1st International Symposium on polarization phenomena in nuclear reactions*. Birkhäuser Basel, 1961.
- [20] H. H. Barschall and W. Haeberli. *Proceedings of the 3rd International Symposium on polarization phenomena in nuclear reactions*. University of Wisconsin Press Madison, 1971. ISBN 0299058905.
- [21] G. G. Ohlsen and Jr. P. W. Keaton. Techniques for measurement of spin 1/2 and spin 1 polarization analyzing tensors. *Nuclear Instruments and Methods*, 109(8):41–59, January 1973. URL <http://www.sciencedirect.com/science/article/pii/0029554X73904503>.
- [22] H. Stein, D. Prasuhn, H. Stockhorst, J. Dietrich, K. Phan, V. Kamerdzhiev, R. Maier, I. N. Meshkov, A. O. Sidorin, and V. V. Parkhomchuk. Current Status of the COSY Electron Cooler (Jülich, Germany). *Atomic Energy*,

- 94(1):24–26, 2003. ISSN 1573-8205. doi: 10.1023/A:1023486402224. URL <http://dx.doi.org/10.1023/A:1023486402224>.
- [23] H. J. Stein, D. Prasuhn, H. Stockhorst, J. Dietrich, K. Fan, V. Kamerdjiev, R. Maier, I. N. Meshkov, A. Sidorin, and V. V. Parkhomchuk. Present Performance of Electron Cooling at COSY-Jülich. In *18th Conference on Charged Particle Accelerators (RUPAC 2002) Ozninsk, Russia, October 1-4, 2002*, 2011. URL <https://inspirehep.net/record/885968/files/arXiv:1101.5963.pdf>.
- [24] D. Prasuhn, J. Dietrich, R. Maier, R. Stassen, H.J. Stein, and H. Stockhorst. Electron and stochastic cooling at {COSY}. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 441(1–2):167 – 174, 2000. ISSN 0168-9002. doi: [http://dx.doi.org/10.1016/S0168-9002\(99\)01128-6](http://dx.doi.org/10.1016/S0168-9002(99)01128-6). URL <http://www.sciencedirect.com/science/article/pii/S0168900299011286>.
- [25] R. Maier. Cooler synchrotron COSY — performance and perspectives. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 390(1–2):1 – 8, 1997. ISSN 0168-9002. doi: [http://dx.doi.org/10.1016/S0168-9002\(97\)00324-0](http://dx.doi.org/10.1016/S0168-9002(97)00324-0). URL <http://www.sciencedirect.com/science/article/pii/S0168900297003240>.
- [26] P. D. Eversheim, M. Altmeier, O. Felden, R. Gebel, M. Krekel, M. Schak, W. Kretschmer, A. Glombik, K. Mümmeler, P. Nebert, G. Suft, R. Weidmann, H. Paetzgen, Schieck, M. Eggert, S. Lemaître, H. Patberg, R. Reckenfelderbäumer, and C. Schneider. Status of the polarized source for the cooler synchrotron COSY Jülich. *AIP Conference Proceedings*, 293(1):92–96, 1993. doi: <http://dx.doi.org/10.1063/1.45153>. URL <http://scitation.aip.org/content/aip/proceeding/aipcp/10.1063/1.45153>.
- [27] W. Haeberli. Sources of polarized ions. *Annual Review of Nuclear Science*, 17(1):373–426, 1967. doi: 10.1146/annurev.ns.17.120167.002105. URL <http://dx.doi.org/10.1146/annurev.ns.17.120167.002105>.
- [28] O. Felden, R. Gebel, R. Maier, and D. Prasuhn. Negative ion source development at the cooler synchrotron COSY/Jülich. In O. Tarvainen and T. Kalvas, editors, *American Institute of Physics Conference Series*, volume 1515 of *American Institute of Physics Conference Series*, pages 321–330, February 2013. doi: 10.1063/1.4792800.

- [29] R. Weidmann, A. Glombik, H. Meyer, W. Kretschmer, M. Altmeier, P. D. Eversheim, O. Felden, R. Gebel, M. Glende, M. Eggert, S. Lemaitre, R. Reckenfelderbäumer, and H. Paetz gen. Schieck. The polarized ion source for COSY. *Review of Scientific Instruments*, 67(3):1357–1358, 1996. doi: <http://dx.doi.org/10.1063/1.1146665>. URL <http://scitation.aip.org/content/aip/journal/rsi/67/3/10.1063/1.1146665>.
- [30] R. Gebel, J. M. Felden, M. Glende, R. Maier, P. von Rossen, and P. D. Eversheim. New developments at the polarized ion source of COSY-Jülich. In *International Workshop on Polarized Ion Sources and Targets ; PST 2001*, 2001. URL <http://juser.fz-juelich.de/record/37443>. Record converted from VDB: 12.11.2012.
- [31] D. Chiladze. *Polarised Charge-Exchange Reaction $dp \rightarrow (pp)n$ Studies at the ANKE-COSY Spectrometer*. PhD thesis, Tbilisi State University, 2008. URL collaborations.fz-juelich.de/ikp/anke/theses/D_Chiladze_PhD.pdf.
- [32] V. S. Morozov, A. D. Krisch, M. A. Leonova, R. S. Raymond, D. W. Sivers, V. K. Wong, R. Gebel, A. Lehrach, B. Lorentz, R. Maier, D. Prasuhn, A. Schnase, H. Stockhorst, D. Eversheim, F. Hinterberger, H. Rohd-jeß, K. Ulbrich, and K. Yonehara. Spin manipulating stored 1.85 GeV/c vector and tensor polarized spin-1 bosons. *Phys. Rev. ST Accel. Beams*, 8:061001, Jun 2005. doi: 10.1103/PhysRevSTAB.8.061001. URL <http://link.aps.org/doi/10.1103/PhysRevSTAB.8.061001>.
- [33] A. Lehrach, U. Bechstedt, J. Dietrich, R. Gebel, B. Lorentz, et al. Acceleration of polarized protons and deuterons at COSY. *AIP Conf.Proc.*, 675(1):153–165, 2003. doi: <http://dx.doi.org/10.1063/1.1607127>. URL <http://scitation.aip.org/content/aip/proceeding/aipcp/10.1063/1.1607127>.
- [34] E. Weise. *Anregungsfunktionen der Analysierstärke der elastischen Proton-Proton-Streuung*. PhD thesis, University of Bonn, 2000. URL <http://edda.hiskp.uni-bonn.de/dipldiss/diss.EW.ps.gz>.
- [35] S. Barsov, U. Bechstedt, W. Bothe, N. Bongers, G. Borchert, W. Borgs, W. Bräutigam, M. Büscher, W. Cassing, V. Chernyshev, B. Chiladze, J. Dietrich, M. Drochner, S. Dymov, W. Erven, R. Esser, A. Franzen, Ye. Golubeva, D. Gotta, T. Grande, D. Grzonka, A. Hardt, M. Hartmann, V. Hejny, L.v. Horn, L. Jarczyk, H. Junghans, A. Kacharava, B. Kamys, A. Khoukaz, T. Kirchner, F. Klehr, W. Klein, H.R. Koch, V.I. Komarov, L. Kondratyuk,

- V Koptev, S Kopyto, R Krause, P Kravtsov, V Kruglov, P Kulesa, A Kulikov, N Lang, N Langenhagen, A Lepges, J Ley, R Maier, S Martin, G Macharashvili, S Merzliakov, K Meyer, S Mikirtychiants, H Müller, P Munhofen, A Mussgiller, M Nekipelov, V Nelyubin, M Nioradze, H Ohm, A Petrus, D Prasuhn, B Prietzschk, H.J Probst, K Pysz, F Rathmann, B Rimarzig, Z Rudy, R Santo, H Paetz gen. Schieck, R Schleichert, A Schneider, Chr Schneider, H Schneider, U Schwarz, H Seyfarth, A Sibirtsev, U Sieling, K Sistemich, A Selikov, H Stechemesser, H.J Stein, A Strzalkowski, K.-H Watzlawik, P Wüstner, S Yashenko, B Zalikhanov, N Zhuravlev, K Zvoll, I Zychor, O.W B. Schult, and H Ströher. ANKE, a new facility for medium energy hadron physics at COSY-Jülich. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 462(3):364 – 381, 2001. ISSN 0168-9002. doi: [http://dx.doi.org/10.1016/S0168-9002\(00\)01147-5](http://dx.doi.org/10.1016/S0168-9002(00)01147-5). URL <http://www.sciencedirect.com/science/article/pii/S0168900200011475>.
- [36] A. Khoukaz, T. Lister, C. Quentmeier, R. Santo, and C. Thomas. Systematic studies on hydrogen cluster beam production. *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics*, 5(2):275–281, 1999. ISSN 1434-6060. doi: 10.1007/PL00021595. URL <http://dx.doi.org/10.1007/PL00021595>.
- [37] S. Dymov, W. Erven, A. Kacharava, R. Koch, V. Komarov, A. Kulikov, V. Kurbatov, G. Macharashvili, H. Ohm, A. Ptrus, F. Rathmann, H. Seyfarth, H. Ströher, S. Yaschenko, B. Zalikhanov, and K. Zvoll. The forward detector of the ANKE spectrometer. Tracking system and its use in data analysis. *Particles and Nuclei, Letters*, 2:40, 2004. URL http://wwwinfo.jinr.ru/publish/Pepan_letters/panl_2_2004/2_2004_05_dym.pdf.
- [38] S. Dymov. *Investigation of the deuteron breakup by protons of 0.6-0.9 GeV with emission of a forward proton pair*. PhD thesis, JINR, 2007. URL http://collaborations.fz-juelich.de/ikp/anke/theses/S.Dymov_PhD.pdf.
- [39] D. Mchedlishvili. *Studies of the neutron-proton charge-exchange amplitudes at COSY using the ANKE spectrometer*. PhD thesis, Tbilisi State University, 2013. URL collaborations.fz-juelich.de/ikp/anke/theses/DM-thesis_13.09.2013.pdf.
- [40] R. Schleichert, T. Krings, S. Merzliakov, A. Mussgiller, and D. Protic. A self-triggering silicon-tracking telescope for spectator proton detection. *Nuclear*

- Science, IEEE Transactions on*, 50(3):301–306, June 2003. ISSN 0018-9499. doi: 10.1109/TNS.2003.812431.
- [41] G. Macharashvili. The software development for the silicon detector data analysis at ANKE-COSY. Technical Note. URL <http://apps.fz-juelich.de/pax/paxwiki/images/6/6c/TechNote15.pdf>.
- [42] G. Macharashvili. A_y measurement in pp elastic scattering. Systematic uncertainties. Technical Note, 2013.
- [43] G. Macharashvili. A neural network concept for event reconstruction at ANKE. Technical Note, 2000.
- [44] S. Dymov. A_y in pp measured with forward detector. Technical Note, 2014.
- [45] S. Yaschenko, S. Dymov, A. Kacharava, V. Komarov, G. Macharashvili, F. Rathmann, S. Barsov, R. Gebel, M. Hartmann, A. Khoukaz, P. Kullessa, A. Kulikov, V. Kurbatov, N. Lang, I. Lehmann, B. Lorentz, T. Mersmann, S. Merzliakov, S. Mikirtytchians, A. Mussgiller, M. Nioradze, H. Ohm, D. Prasuhn, R. Schleichert, H. Seyfarth, E. Steffens, H. J. Stein, H. Ströher, Yu. Uzikov, B. Zalikhanov, and N. Zhuravlev. Measurement of the analyzing power in $\vec{p} \, d \rightarrow (pp)n$ with a fast forward 1S_0 proton pair. *Phys. Rev. Lett.*, 94:072304, Feb 2005. doi: 10.1103/PhysRevLett.94.072304. URL <http://link.aps.org/doi/10.1103/PhysRevLett.94.072304>.
- [46] M. W. McNaughton, P. R. Bevington, H. B. Willard, E. Winkelmann, E. P. Chamberlin, F. H. Cverna, N. S. P. King, and H. Willmes. Polarization analyzing power $A(\theta)$ in pp elastic scattering at 796 MeV. *Phys. Rev. C*, 23:1128–1133, Mar 1981. doi: 10.1103/PhysRevC.23.1128. URL <http://link.aps.org/doi/10.1103/PhysRevC.23.1128>.
- [47] F. Irom, G. J. Igo, J. B. McClelland, and C. A. Whitten. Elastic $p - p$ scattering at 796 MeV in the coulomb-nuclear interference region. *Phys. Rev. C*, 25:373–386, Jan 1982. doi: 10.1103/PhysRevC.25.373. URL <http://link.aps.org/doi/10.1103/PhysRevC.25.373>.
- [48] P. R. Bevington, M. W. McNaughton, H. B. Willard, H. W. Baer, E. Winkelmann, F. Cverna, E. P. Chamberlin, N. S. P. King, R. R. Stevens, H. Wilmes, and M. A. Schardt. Polarization analyzing power $A_y(\theta)$ in pp elastic scattering at 643, 787, and 796 MeV. *Phys. Rev. Lett.*, 41:384–387, Aug 1978. doi: 10.1103/PhysRevLett.41.384. URL <http://link.aps.org/doi/10.1103/PhysRevLett.41.384>.

- [49] C.E. Allgower, J. Ball, M. Beddo, Y. Bedfer, A. Boutefnouchet, J. Bystricky, P.-A. Chamouard, Ph. Demierre, J.-M. Fontaine, V. Ghazikhanian, D. Grossnick, R. Hess, Z. Janout, Z.F. Janout, V.A. Kalinnikov, T.E. Kasprzyk, B.A. Khachaturov, R. Kunne, F. Lehar, A. de Lesquen, D. Lopiano, V.N. Matafonov, I.L. Oisarev, A.A. Popov, A.N. Prokofiev, D. Rapin, J.-L. Sans, H.M. Spinka, A. Teglia, Yu.A. Usov, V.V. Vikhrov, B. Vuaridel, C.A. Whitten, and A.A. Zhdanov. The pp elastic scattering analyzing power measured with the polarized beam and the unpolarized target between 1.98 and 2.80 GeV. *Nuclear Physics A*, 637(2):231 – 242, 1998. ISSN 0375-9474. doi: [http://dx.doi.org/10.1016/S0375-9474\(98\)00216-4](http://dx.doi.org/10.1016/S0375-9474(98)00216-4). URL <http://www.sciencedirect.com/science/article/pii/S0375947498002164>.
- [50] Z. Bagdasarian, D. Chiladze, S. Dymov, A. Kacharava, G. Macharashvili, S. Barsov, R. Gebel, B. Gou, M. Hartmann, I. Keshelashvili, A. Khoukaz, P. Kulesa, A. Kulikov, A. Lehrach, N. Lomidze, B. Lorentz, R. Maier, D. Mchedlishvili, S. Merzliakov, S. Mikirtychyants, M. Nioradze, H. Ohm, M. Papenbrock, D. Prasuhn, F. Rathmann, V. Serdyuk, V. Shmakova, R. Stassen, H. Stockhorst, I.I. Strakovsky, H. Ströher, M. Tabidze, A. Täschner, S. Trusov, D. Tsirkov, Yu. Uzikov, Yu. Valdau, C. Wilkin, and R.L. Workman. Measurement of the analysing power in proton–proton elastic scattering at small angles. *Physics Letters B*, 739:152 – 156, 2014. ISSN 0370-2693. doi: <http://dx.doi.org/10.1016/j.physletb.2014.10.054>. URL <http://www.sciencedirect.com/science/article/pii/S0370269314007874>.
- [51] Private Communications with R. Workman and I. Strakovsky of SAID group.
- [52] H. J. Stein, M. Hartmann, I. Keshelashvili, Y. Maeda, C. Wilkin, S. Dymov, A. Kacharava, A. Khoukaz, B. Lorentz, R. Maier, T. Mersmann, S. Mikirtychyants, D. Prasuhn, R. Stassen, H. Stockhorst, H. Ströher, Yu. Valdau, and P. Wüstner. Determination of target thickness and luminosity from beam energy losses. *Phys. Rev. ST Accel. Beams*, 11:052801, May 2008. doi: 10.1103/PhysRevSTAB.11.052801. URL <http://link.aps.org/doi/10.1103/PhysRevSTAB.11.052801>.
- [53] Kirsten Zapfe, W. Brückner, H.-G. Gaul, M. Grieser, M.T. Lin, Z. Moroz, B. Povh, M. Rall, B. Stechert, E. Steffens, J. Stenger, F. Stock, J. Tonhäuser, Ch. Montag, F. Rathmann, D. Fick, B. Braun, G. Graw, and W. Haeberli. Detailed studies of a high-density polarized hydrogen gas target for storage rings. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated*

- Equipment*, 368(2):293 – 306, 1996. ISSN 0168-9002. doi: [http://dx.doi.org/10.1016/0168-9002\(95\)00608-7](http://dx.doi.org/10.1016/0168-9002(95)00608-7). URL <http://www.sciencedirect.com/science/article/pii/0168900295006087>.
- [54] National Institute of Standards and Technologies. Physical Measurement Laboratory (NIST-PML) database. URL <http://nist.gov/pml/data/star/>.
- [55] W. Schottky. Zur theorie des elektronenrauschens in mehrgitterröhren. *Annalen der Physik*, 424(1-2):195–204, 1938. ISSN 1521-3889. doi: 10.1002/andp.19384240122. URL <http://dx.doi.org/10.1002/andp.19384240122>.
- [56] S. Dymov. The proton-proton elastic scattering count rates from the ANKE Forward Detector: Systematic uncertainties.
- [57] D. Mchedlishvili, D. Chiladze, S. Dymov, Z. Bagdasarian, S. Barsov, R. Gebel, B. Gou, M. Hartmann, A. Kacharava, I. Keshelashvili, A. Khoukaz, P. Kulesa, A. Kulikov, A. Lehrach, N. Lomidze, B. Lorentz, R. Maier, G. Macharashvili, S. Merzliakov, S. Mikirtychyants, M. Nioradze, H. Ohm, D. Prasuhn, F. Rathmann, V. Serdyuk, D. Schroer, V. Shmakova, R. Stassen, H.J. Stein, H. Stockhorst, I.I. Strakovsky, H. Ströher, M. Tabidze, A. Täschner, S. Trusov, D. Tsirkov, Yu. Uzikov, Yu. Valdau, C. Wilkin, R.L. Workman, and P. Wüstner. Measurement of the absolute differential cross section of proton–proton elastic scattering at small angles. *Physics Letters B*, 755:92 – 96, 2016. ISSN 0370-2693. doi: <http://dx.doi.org/10.1016/j.physletb.2016.01.066>. URL <http://www.sciencedirect.com/science/article/pii/S037026931600085X>.
- [58] A.V. Dobrovolsky, A.V. Khanzadeev, G.A. Korolev, E.M. Maev, V.I. Medvedev, G.L. Sokolov, N.K. Terentyev, Y. Terrien, G.N. Velichko, A.A. Vorobyov, and Yu.K. Zalite. Small angle pp scattering at energies from 650 to 1000 MeV. *Nuclear Physics B*, 214(1):1 – 20, 1983. ISSN 0550-3213. doi: [http://dx.doi.org/10.1016/0550-3213\(83\)90163-3](http://dx.doi.org/10.1016/0550-3213(83)90163-3). URL <http://www.sciencedirect.com/science/article/pii/0550321383901633>.
- [59] B.H. Silverman, J.C. Lugol, J. Saudinos, Y. Terrien, F. Wellers, A.V. Dobrovolsky, A.V. Khanzadeev, G.A. Korolev, G.E. Petrov, E.M. Spiridenkov, and A.A. Vorobyov. Differential cross sections and analyzing powers for small angle neutron-proton elastic scattering between 378 and 1135 MeV. *Nuclear Physics A*, 499(4):763 – 788, 1989. ISSN 0375-9474. doi: [http://dx.doi.org/10.1016/0375-9474\(89\)90062-6](http://dx.doi.org/10.1016/0375-9474(89)90062-6). URL <http://www.sciencedirect.com/science/article/pii/0375947489900626>.

- [60] A. V. Dobrovolsky, A. V. Khanzadeev, G. A. Korolev, G. N. Velichko, A. A. Vorobyov, J. Saudinos, B. H. Silverman, Y. Terrien, and F. Wellers. Experimental data on elastic pp, np, pd, and p α forward scattering at intermediate energies. Technical Report LENINGRAD-INST-YAD-FIZ-1454. LIYF-1454, Akad. Nauk St. Petersburg. Inst. Yarn. Fiz., St. Petersburg, Nov 1988. URL <https://cds.cern.ch/record/195992>.
- [61] I. Ambats, D. S. Ayres, R. Diebold, A. F. Greene, S. L. Kramer, A. Lesnik, D. R. Rust, C. E. W. Ward, A. B. Wicklund, and D. D. Yovanovitch. Systematic study of $\pi^\pm p$, $K^\pm p$, pp, and $\bar{p}p$ forward elastic scattering from 3 to 6 GeV/c. *Phys. Rev. D*, 9:1179–1209, Mar 1974. doi: 10.1103/PhysRevD.9.1179. URL <http://link.aps.org/doi/10.1103/PhysRevD.9.1179>.
- [62] A precision measurement of pp elastic scattering cross-sections at intermediate energies. *The European Physical Journal A - Hadrons and Nuclei*, 22(1):125–148, 2004. ISSN 1434-6001. doi: 10.1140/epja/i2004-10011-3. URL <http://dx.doi.org/10.1140/epja/i2004-10011-3>.
- [63] W. Grein and P. Kroll. Amplitude analysis of forward nucleon-nucleon scattering. *Nuclear Physics A*, 377(2):505 – 517, 1982. ISSN 0375-9474. doi: [http://dx.doi.org/10.1016/0375-9474\(82\)90052-5](http://dx.doi.org/10.1016/0375-9474(82)90052-5). URL <http://www.sciencedirect.com/science/article/pii/0375947482900525>.
- [64] Richard A. Arndt, L. David Roper, Ronald A. Bryan, Robert B. Clark, Bruce J. VerWest, and Peter Signell. Nucleon-nucleon partial-wave analysis to 1 GeV. *Phys. Rev. D*, 28:97–122, Jul 1983. doi: 10.1103/PhysRevD.28.97. URL <http://link.aps.org/doi/10.1103/PhysRevD.28.97>.
- [65] C. Lechanoine, F. Lehar, F. Perrot, and P. Winternitz. Polarization phenomena in nucleon-nucleon forward scattering. *Il Nuovo Cimento A Series 11*, 56(2):201–228, 1980. ISSN 0369-3546. doi: 10.1007/BF02730164. URL <http://dx.doi.org/10.1007/BF02730164>.
- [66] Pluto. A monte carlo simulation tool for hadronic physics. URL <https://www-hades.gsi.de/?q=pluto>.
- [67] A. Mussgiller. *Identification and Tracking of Low Energy Spectator Protons*. PhD thesis, University of Cologne, 2007. URL collaborations.fz-juelich.de/ikp/anke/theses/A.Mussgiller_PhD.pdf.
- [68] F. Irom, G. J. Igo, J. B. McClelland, C. A. Whitten, and M. Bleszynski. Measurements of small angle elastic $\vec{p}$ -d scattering at 796 MeV using a recoil

- method. *Phys. Rev. C*, 28:2380–2385, Dec 1983. doi: 10.1103/PhysRevC.28.2380. URL <http://link.aps.org/doi/10.1103/PhysRevC.28.2380>.
- [69] M. L. Barlett, G. W. Hoffmann, J. A. McGill, B. Hoistad, L. Ray, R. W. Ferguson, E. C. Milner, J. A. Marshall, J. F. Amann, B. E. Bonner, J. B. McClelland, G. S. Blanpied, and R. A. Arndt. Forward-angle elastic and quasielastic proton-nucleon cross sections and analyzing powers at 0.8 GeV. *Phys. Rev. C*, 27:682–696, Feb 1983. doi: 10.1103/PhysRevC.27.682. URL <http://link.aps.org/doi/10.1103/PhysRevC.27.682>.
- [70] J. Ball, Ph. Chesny, M. Combet, J.M. Fontaine, C.D. Lac, J.L. Sans, J. Bystricky, F. Lehar, A. de Lesquen, M. de Mali, F. Perrot-Kunne, L. van Rossum, A. Ahmidouch, P. Bach, Ph. Demierre, G. Gaillard, R. Hess, R. Kunne, D. Rapin, Ph. Sormani, J.P. Goudour, R. Binz, A. Klett, E. Rössle, H. Schmitt, D. Lopiano, and H. Spinka. Angular dependence of analyzing power in np elastic scattering between 0.312 and 1.10 GeV. *Nuclear Physics A*, 559(4):489 – 510, 1993. ISSN 0375-9474. doi: [http://dx.doi.org/10.1016/0375-9474\(93\)90257-X](http://dx.doi.org/10.1016/0375-9474(93)90257-X). URL <http://www.sciencedirect.com/science/article/pii/037594749390257X>.
- [71] G. Glass, T. S. Bhatia, J. C. Hiebert, R. A. Kenefick, S. Nath, L. C. Northcliffe, K. F. Johnson, H. Spinka, R. Stanek, M. W. Rawool, J. A. Faucett, R. H. Jeppesen, G. E. Tripard, and C. R. Newsom. Analyzing power measurement for forward angle $n - p$ scattering at 790 MeV. *Phys. Rev. C*, 41:2732–2736, Jun 1990. doi: 10.1103/PhysRevC.41.2732. URL <http://link.aps.org/doi/10.1103/PhysRevC.41.2732>.
- [72] G. Glass, T. S. Bhatia, J. C. Hiebert, R. A. Kenefick, S. Nath, L. C. Northcliffe, W. B. Tippens, and J. E. Simmons. Forward angle analyzing power in $p \rightarrow n$ and $p \rightarrow p$ quasifree scattering at 643 and 797 mev. *Phys. Rev. C*, 47:1369–1375, Apr 1993. doi: 10.1103/PhysRevC.47.1369. URL <http://link.aps.org/doi/10.1103/PhysRevC.47.1369>.
- [73] R. Diebold, D. S. Ayres, S. L. Kramer, A. J. Pawlicki, and A. B. Wicklund. Measurement of the proton-neutron elastic-scattering polarization from 2 to 6 GeV/c. *Phys. Rev. Lett.*, 35:632–635, Sep 1975. doi: 10.1103/PhysRevLett.35.632. URL <http://link.aps.org/doi/10.1103/PhysRevLett.35.632>.
- [74] Y. Makdisi, M. L. Marshak, B. Mossberg, E. A. Peterson, K. Ruddick, J. B. Roberts, and R. D. Klem. Analyzing power in large-angle proton-neutron elastic scattering. *Phys. Rev. Lett.*, 45:1529–1533, Nov 1980. doi:

- 10.1103/PhysRevLett.45.1529. URL <http://link.aps.org/doi/10.1103/PhysRevLett.45.1529>.
- [75] S. Nurushev, M. Runtso, and M. Strikhanov. *Introduction to Polarization Physics*, volume 859. Springer-Verlag Berlin Heidelberg, 2013. doi: 10.1007/978-3-642-32163-4. URL <http://www.springer.com/la/book/9783642321627>.
- [76] J. H. Parry, N. E. Booth, G. Conforto, R. J. Esterling, J. Scheid, D. J. Sherden, and A. Yokosawa. Measurements of the polarization in proton-proton elastic scattering from 2.50 to 5.15 GeV/ c . *Phys. Rev. D*, 8:45–63, Jul 1973. doi: 10.1103/PhysRevD.8.45. URL <http://link.aps.org/doi/10.1103/PhysRevD.8.45>.
- [77] S. Barsov, Z. Bagdasarian, D. Chiladze, S. Dymov, and A. Kacharava. Measurements of the pn quasi-free elastic scattering at anke, 2015. URL collaborations.fz-juelich.de/ikp/anke/annual/annual_reports/14/pnQuasi_status_2014.pdf.
- [78] I. Lehmann. ω Meson Production in the $pn \rightarrow d\omega$ Reaction at ANKE. PhD thesis, University of Cologne, 2003. URL collaborations.fz-juelich.de/ikp/anke/theses/I.Lehmann_PhD.pdf.

Erklärung

Ich versichere, dass ich die von mir vorgelegte Dissertation selbständig angefertigt, die benutzten Quellen und Hilfsmittel vollständig angegeben und die Stellen der Arbeit – einschliesslich Tabellen, Karten und Abbildungen –, die anderen Werken im Wortlaut oder dem Sinn nach entnommen sind, in jedem Einzelfall als Entlehnung kenntlich gemacht habe; dass diese Dissertation abgesehen von der Staatlichen Iwane-Dschawachischwili-Universität Tiflis noch keiner anderen Fakultät oder Universität zur Prüfung vorgelegen hat; dass sie – abgesehen von unten angegebenen Teilpublikationen – noch nicht veröffentlicht worden ist, sowie, dass ich eine solche Veröffentlichung vor Abschluss des Promotionsverfahrens nicht vornehmen werde. Die Bestimmungen der Promotionsordnung sind mir bekannt. Die von mir vorgelegte Dissertation ist von Prof. Dr. Dr. h.c. Hans Ströher und Dr. Nodar Lomidze betreut worden.

Publikationen

mit Daten aus der vorliegenden Dissertation

- [1] Z. Bagdasarian, D. Chiladze, S. Dymov, A. Kacharava, G. Macharashvili, S. Barsov, R. Gebel, B. Gou, M. Hartmann, I. Keshelashvili, A. Khoukaz, P. Kulesa, A. Kulikov, A. Lehrach, N. Lomidze, B. Lorentz, R. Maier, D. Mchedlishvili, S. Merzliakov, S. Mikirtychyants, M. Nioradze, H. Ohm, M. Papenbrock, D. Prasuhn, F. Rathmann, V. Serdyuk, V. Shmakova, R. Stassen, H. Stockhorst, I.I. Strakovsky, H. Ströher, M. Tabidze, A. Täschner, S. Trusov, D. Tsirkov, Yu. Uzikov, Yu. Valdau, C. Wilkin, R.L. Workman, **Measurement of the analysing power in proton–proton elastic scattering at small angles**, Physics Letters B, Volume 739 (2014) 152, ISSN 0370-2693 doi:<http://dx.doi.org/10.1016/j.physletb.2014.10.054>
- [2] Z. Bagdasarian for the ANKE collaboration, **Nucleon-nucleon scattering at small angles measured at ANKE-COSY** EPJ Web of Conferences Volume 113 (2016) 04023. Proceedings of the 21st International Conference on Few-body Problems in Physics, Chicago, USA. doi:<http://dx.doi.org/10.1051/epjconf/201611304023>
- [3] D. Mchedlishvili, D. Chiladze, S. Dymov, Z. Bagdasarian, S. Barsov, R. Gebel, B. Gou, M. Hartmann, A. Kacharava, I. Keshelashvili, A. Khoukaz, P. Kulesa, A. Kulikov, A. Lehrach, N. Lomidze, B. Lorentz, R. Maier, G. Macharashvili, S. Merzliakov, S. Mikirtychyants, M. Nioradze, H. Ohm, D. Prasuhn, F. Rathmann, V. Serdyuk, D. Schroer, V. Shmakova, R. Stassen, H.J. Stein, H. Stockhorst, I.I. Strakovsky, H. Ströher, M. Tabidze, A. Täschner,

S. Trusov, D. Tsirkov, Yu. Uzikov, Yu. Valdau, C. Wilkin, R.L. Workman, P. Wüstner,

Measurement of the absolute differential cross section of proton–proton elastic scattering at small angles, Physics Letters B, Volume 755 (2016) 92, ISSN 0370-2693, doi:<http://dx.doi.org/10.1016/j.physletb.2016.01.066>

[4] Z. Bagdasarian for the ANKE collaboration,

Measurement of the analysing power in proton–proton elastic scattering, Proceedings of Science (2015) 53rd International Winter Meeting on Nuclear Physics, Bormio, Italy. http://pos.sissa.it/archive/conferences/238/014/Bormio2015_014.pdf

Lebenslauf

Persönliche Daten

Name:	Zara Bagdasarian
Geburtsdatum:	04.10.1989
Geburstort:	Tiflis, Georgien
Staatsangehörigkeit:	Georgisch
Familienstand:	Verheiratet

Promotionsstudium

seit November 2012	Staatlichen Iwane-Dschawachischwili-Universität Tiflis und Universität zu Köln (Cotutelle Verfahren)
Doktorarbeit	Measurement of Nucleon-Nucleon Elastic Scattering at Small Angles using the ANKE spectrometer

Studium

Oktober 2010 - Juli 2012	Master der Physik der Staatlichen Iwane-Dschawachischwili-Universität Tiflis, Georgien Gesamtnote: "mit Auszeichnung"
Masterarbeit	Beam Polarimetry Studies for Spin-Filtering Experiment at COSY, Jülich
Oktober 2006 - Juli 2010	Bachelor der Physik der Staatlichen Iwane-Dschawachischwili-Universität Tiflis, Georgien Gesamtnote: "mit Auszeichnung"

Schulbildung

September 1996 - Juni 2006	Schule #9, Tiflis, Georgien Gesamtnote: "mit Auszeichnung"
----------------------------	---